

DESCRIPTION

MATERIALS AND METHODS FOR CONTROLLING PESTS

5

Government Support

The subject invention was made with funding from the National Institutes of Health (Grant No. AI 030464). Accordingly, the government may have certain rights in this invention.

Background of Invention

10

A longstanding worldwide demand exists for new, effective, environmentally friendly, and safe means to control pests that damage agriculture or serve as disease vectors. Agriculture costs incurred by pests exceed billions of dollars annually in decreased crop yields, reduced crop quality, increased harvesting costs, pesticide application costs, and negative ecological impact. In addition to agriculture pests, many blood-feeding insects are vectors for pathogenic microorganisms that threaten human and animal health, or are annoying at the least. As in the case of agriculture pests, direct and intangible costs incurred by blood-feeding pests concern pesticide safety hazards to humans and animals, bioaccumulation and environmental incompatibility, and synthesis and application costs.

15

20

25

30

Almost all field crops, nursery and horticulture plants, and commercial farming areas are susceptible to attack by one or more pests. Particularly problematic are Coleopteran and Lepidopteran pests. An example of a Lepidopteran pest is the hornworm larva of *Manduca sexta*, and an example of a Coleopteran pest is the Colorado potato beetle, *Leptinotarsa decemlineata*. Vegetable and cole crops, lentils, leafy vegetables, melons, peppers, potatoes and related tubers, tomatoes, cucumbers and related vine crops, as well as a variety of spices are sensitive to infestation by one or more pests including loopers, armyworms, moth larvae, budworms, webworms, earworms, leafeaters, borers, cloverworms, melonworms, leafrollers, various caterpillars, fruitworms, hornworms, and pinworms. Likewise, pasture and hay crops such as alfalfa, pasture and forage grasses and silage are often attacked by a variety of pests including armyworms, alfalfa caterpillars, European skipper, a variety of loopers and webworms, as well as yellowstriped armyworms.

Fruit (including citrus), nut, and vine crops are susceptible to attack by a variety of pests, including sphinx moth larvae, cutworms, skippers, fireworms, leafrollers, cankerworms, fruitworms, girdlers, webworms, leaffolders, skeletonizers, shuckworms, hornworms, loopers, orangeworms, tortrix, twig borers, casebearers, spanworms, budworms, budmoths, and a variety of caterpillars and armyworms.

Field crops are targets for infestation by insects including armyworm, asian and other corn borers, a variety of moth and caterpillar larvae, bollworms, loopers, rootworms, leaf perforators, cloverworms, headworms, cabbageworms, leafrollers, podworms, cutworms, budworms, hornworms, and the like. Pests also frequently feed upon bedding plants, flowers, ornamentals, vegetables, container stock, forests, fruit, ornamental, shrubs and other nursery stock. Even turf grasses are attacked by a variety of pests including armyworms and sod webworms.

For the past 50 years growers, health officials, and the public have depended on chemical pesticides for controlling a variety of pests. However, environmental experts, health officials, and the public have become concerned about the amount of residual chemicals found in food, ground water, and elsewhere in the environment. Regulatory agencies around the world are restricting and/or banning the uses of many synthetic pesticides, particularly those that are persistent in the environment and that enter the food chain. Stringent new restrictions on the use of pesticides and the elimination of some effective pesticides from the market place could limit economical and effective options for controlling costly pests. Some synthetic chemical pesticides can poison the soil and underlying aquifers, pollute surface waters as a result of runoff, and destroy non-target life forms. These synthetic chemical pest control agents have the further disadvantage of presenting public safety hazards when they are applied in areas where pets, farm animals, or children may come into contact with them. They can also pose health hazards to the people applying them, especially if the proper application techniques are not followed.

Because crops of commercial interest are often the targets of pests, environmentally sensitive methods for controlling or eradicating pest infestations are desirable in many instances. This is particularly true for farmers, nurserymen, growers, and commercial and residential areas which seek to control pest populations using environmentally friendly compositions.

The most widely used environmentally friendly pesticidal formulations developed in recent years have been microbial pesticides derived from the bacterium *Bacillus thuringiensis* ("B.t."). *B. thuringiensis* is a Gram-positive bacterium that produces proteins which are toxic to certain orders and species of pests. Many different strains of *B. thuringiensis* have been shown to produce insecticidal proteins. Compositions including *B. thuringiensis* strains which produce insecticidal proteins have been commercially-available and used as environmentally-acceptable insecticides because they are toxic to specific target pests, but are harmless to plants and other non-target organisms. The specificity of these toxins is often strain-specific, with certain toxins being active against a relatively narrow spectrum of pests. Indeed, many *B.t.* toxins have been identified that are active only against particular insect orders (e.g., dipterans, hymenopterans, coleopterans, etc.). This limitation prevents the use of a single *B.t.* endotoxin composition as a broad-range pesticide.

Crop pests are not the only targets for which an environmentally friendly and safe pesticide would be highly desirable. Many blood-feeding pests are known to prey on humans and animals, and many pests are vectors for pathogenic microorganisms that threaten human and animal health, including commercially important livestock, pets and other animals. The order Diptera contains a large number of blood-ingesting and disease-carrying pests, including, for example, mosquitoes, black flies, no-see-ums (punkies), horse flies, deer flies and tsetse flies. Various species of mosquitoes, for example, transmit diseases caused by viruses, and many are vectors for disease-causing nematodes and protozoa. Mosquitoes of the genus *Anopheles* transmit *Plasmodium*, the protozoan that causes malaria. The mosquito species *Aedes aegypti* transmits an arbovirus that causes yellow fever in humans. Other viruses transmitted by *Aedes* species include the causative agents of dengue fever, eastern and western encephalitis, Venezuelan equine encephalitis, St. Louis encephalitis, chikungunya, oroponehe and bunyarnidera. The genus *Culex*, which includes the common house mosquito *C. pipiens*, is implicated in the transmission of various forms of encephalitis, filarial worms, and West Nile virus. *Trypanasoma cruzi*, the causative agent of Chagas disease, is transmitted by various species of blood ingesting Triatominae bugs. Tsetse flies (*Glossina spp.*) transmit African trypanosomal diseases of humans and cattle. Other diseases are transmitted by various blood-ingesting pest species.

Various pesticides have been employed in efforts to control or eradicate populations of disease-transmitting pests. For example, DDT, a chlorinated hydrocarbon, has been used in attempts to eradicate malaria-bearing mosquitoes throughout the world. Other examples of chlorinated hydrocarbons are BHC, lindane, chlorobenzilate, methoxychlor, and the cyclodienes (e.g., aldrin, dieldrin, chlordane, heptachlor, and endrin). The long-term stability of many of these pesticides and their tendency to bioaccumulate render them particularly dangerous to many non-target organisms.

In addition to environmental concerns, another major problem associated with conventional chemical control practices is the capability of many species to develop pesticide resistance. Resistance results from the selection of naturally occurring mutants possessing biochemical, physiological or behavioral factors that enable the pests to tolerate the pesticide when it is applied.

There is clearly a longstanding need in the art for pesticidal compounds that reduce or eliminate direct and/or indirect threats to human health posed by presently available pesticides, that are environmentally compatible and safe, are not toxic to non-pest organisms, and have a reduced tendency to bioaccumulate.

Approaches to pesticide development are lacking that involve specifically disrupting key pest metabolic regulatory processes, notably membrane transporter or channel proteins as targets. The development of such methodologies could provide safer, environmentally friendly alternatives to conventional commercially used pesticides, and provide more economical means for suppressing or eradicating target pest populations. The formulation of pesticidal compositions that are non-toxic to animals and to humans would greatly enhance the present methods available for killing pests, and would provide alternative strategies for environmentally responsible pest management.

Membrane transporter proteins and ion channel proteins serve critical roles in maintaining organic solute and ionic metabolic, thermodynamic, and electrical events in all cells. In both eukaryotes and prokaryotes these proteins affect electrochemical gradients of a wide variety of metabolic molecules and electrolytes, including amino acids and related metabolites as well as H^+ , OH^- , Na^+ , K^+ , Cl^- , and carbonate ions (Gerencser and Stevens, 1994; Stevens, B.R. 2001. "Theory and methods in nutrient membrane transport." In: *Surgical Research*. pp. 845-

856. W.W. Souba and D.W. Wilmore, eds., Academic Press, San Diego). Molecular cloning studies have identified several subfamilies of organic solute transporters and ion channels (Griffith and Sansom, book 1998).

Organic solute transporters and ion channels are commonly defined by their substrate selectivity within polypeptide superfamilies. For cloned or native secondary active transporters, it is generally assumed that cell membranes utilize ion and organic molecule electrochemical gradients to aid in exchanging these solutes between the cell interior and extracellular environment (Gerencser, G.A. and B.R. Stevens, 1994, *J. Exper. Biol.* 196:59-75; Stevens, B.R. 1999, Digestion and Absorption of Protein. In: Biochemical and Physiological Aspects of Human Nutrition. pp. 107-123, M.H. Stipanuk, ed., W.B. Saunders Co., Philadelphia.; <http://www.biology.ucsd.edu/~msaier/transport/>). In the 'prototypical' transporter, organic solutes that can be moved across cell membranes by uniport, hetero- or homo-exchange, and/or uptake can be activated by ions, and/or thermodynamically cotransported with ions (cf refs in Quick, M. and B.R. Stevens, 2001, *J. Biol. Chem.* 276(36):33413-33418; Griffith, J.K. and C.E. Sansom, 1998, In: The Transporter Facts Book, Academic Press, San Diego, pp. 500; <http://www.biology.ucsd.edu/~msaier/transport/>). Ion channels, on the other hand, are typically distinct from organic solute transporters, are selective in their conducting ion species, and may be gated by organic ligands (Hille, B, 2001, *Ionic channels of excitable membranes*, 3rd Edition, Sinauer Associates, Inc., Sunderland, Mass., pp. 814; <http://www.biology.ucsd.edu/~msaier/transport/>).

Manduca sexta is a major crop pest whose larval stage, commonly known as tobacco and tomato hornworms, rapidly attack and defoliate tobacco and tomato plants; the large fifth instar larvae are especially damaging. Other vegetable crops such as peppers, eggplant, and potatoes also can be affected. Tobacco and tomato hornworms rapidly grow and gain weight as they progress from the first instar stage (about 6.7 mm) through the fifth instar (about 81.3 mm) over a period of about 20 days. The sated larvae then drop to the soil to pupate, and eventually emerge as adult moths. The moths lay eggs, which develop into larvae, and the life cycle continues, thereby sustaining crop damage. Killing the larvae prevents immediate crop damage and prevents or reduces future damage by interrupting the life cycle.

The midgut region of *M. sexta* larvae displays compartments with the property of high concentrations of K^+ as well as Na^+ in an alkaline fluid ($\sim pH$ 10), with trans-epithelial potentials ~ 250 mV (Harvey *et al.*, 1999; Harvey and Wieczorek, 1997). Epithelial cells of this region transport a variety of nutrients, including nutrient amino acids and electrolytes, as demonstrated by *in vitro* isolated membrane vesicle uptake studies. In place of a Na^+/K^+ -ATPase typically found in cells, this tissue instead possesses a proton translocating V-ATPase (Graf *et al.*, 1992; Merzendorfer *et al.*, 1997) which energizes the cell membranes for secretion and absorption of K^+ and Na^+ ions, and establishment of a large pH gradient. A K^+ -activated leucine-preferring transporter (KAAT1) has been identified from the hornworm midgut (Castagna *et al.*, 1998, *Proc Natl. Acad. Sci. USA* 95:5395-5400), and a GABA (gamma aminobutyric acid) transporter has been cloned from an *M. sexta* embryo cDNA library (Mbungu *et al.*, 1995).

CAATCH1 (Cation-Amino Acid Transporter/CHannel) is a recently cloned insect membrane protein initially cloned from *Manduca sexta*; CAATCH1 exhibiting a unique polypeptide and nucleotide sequence related to, but different from, mammalian Na^+ -, Cl^- -coupled neurotransmitter transporters (Feldman *et al.*, 2000, *J. Biol. Chem.* 275:24518–24526). Utilizing a unique PCR-based strategy, the gene encoding CAATCH1 was cloned (Feldman *et al.*, 2000, *supra*) from a cDNA library in LambdaZap plasmids, obtained from the digestive midgut of *Manduca sexta* larvae.

The unanticipated and novel biochemical, physiological, and molecular properties of CAATCH1 indicated that it is a multi-function protein that mediates amino acid uptake in a manner that is thermodynamically uncoupled from ion electrochemical potentials, and furthermore that CAATCH1 simultaneously functions as an amino acid-modulated gated alkali cation channel (Quick, M., and B.R. Stevens, 2001, "Amino acid transporter CAATCH1 is also an amino acid-gated cation channel". *J. Biol. Chem.* 276: 33413-33418) serving at least Na^+ and K^+ . Radiotracer and electrophysiology experiments with functional CAATCH1 polypeptide expressed from the full length CAATCH1 cDNA demonstrated direct amino acid ligand-protein interactions, and indicated that binding by different amino acid substrates differentially affects the conformational states of CAATCH1 (Quick, M. and B.R. Stevens, 2001, *J. Biol. Chem.* 276:33413-33418). Notably, L-methionine binding to CAATCH1 *in situ* in biomembranes in the presence of Na^+ perturbs the charge-voltage relation with a high affinity binding constant,

affecting transient currents due to CAATCH1-associated charge transfer across the membrane dielectric field. Furthermore, CAATCH1-associated voltage-dependent amino acid-elicited steady state inward cation currents are blocked by methionine, and indeed methionine reversed charge movements via CAATCH1 expressed in cell membranes, even though radiotracer methionine influx is catalyzed by CAATCH1 (Quick, M. and B.R. Stevens, 2001, *J. Biol. Chem.* 276:33413-33418). In insects, CAATCH1 likely plays a broader role than that of a 'classical' transporter or channel; as a multifunction protein CAATCH1 is likely a key protein in electrolyte and organic solute homeostasis of certain insects (Feldman *et al*, 2000, *J. Biol. Chem.* 275(32):24518-24526)

Many pests—including mosquitoes, Lepidopterans, and Coleopterans—possess an alkaline pH midgut, and share some similar physiological mechanisms that occur within this unusual milieu (Nation, J., 2001, *In: Insect Physiology and Biochemistry*, CRC Press, Boca Raton, pp 496). Mosquito larvae possess such an alkaline midgut, and adjust free amino acid concentrations in their hemolymph and extracellular compartments in direct response to existence of foreign factors in the gut (e.g., *B.t.* δ -endotoxin) or the salinity of their habitat (Bounias, M. *et al.*, 1989, *J. Invertebr Pathol.* 54:16-22). Notably, one standout free amino acid, L-proline, accumulates 4-fold during the normal course of *Aedes aegypti* larval development, and in *Culex spp.* L-proline accumulation can exceed 50-fold (up to 70 mM) when larvae are stimulated by Na^+ in their feeding pools (Bounias, M. *et al.*, 1989, *supra*; Patrick, M.L. and T.J. Bradley, 2000, *J. Exp. Biol.* 203:831-839; Chaput, R.L. and J.N. Liles, 1969, *Ann. Entomol. Soc. Am.* 62:742-747). This proline is likely utilized as an energy source and for osmoregulation (Patrick, M.L. and T.J. Bradley, 2000, *J. Exp. Biol.* 203:831-839; Bounias *et al.*, 1989, *J. Invertebr Pathol.* 54:16-22). In contrast, free L-methionine has the distinction of virtually the lowest measurable concentration (≤ 0.001 mM) of any of the free amino acids in mosquito larvae. Virtually all methionine existing in the free amino acid state in larvae (Dadd, R.H., 1973, *Ann Rev Entomol* 18:381-420) is metabolically shunted and sequestered into so-called methionine-rich hexamerin proteins (Korochkina *et al.*, 1997, *Insect Biochem Mol Biol.* 27:813-824) that are stored for post-larval developmental events. In its role as a nutrient transporter, the CAATCH1 protein has been shown to be primarily responsible for proline uptake (Feldman *et al.*, 2000, *J. Biol. Chem.* 275(32):24518-24526), while in the presence of extremely low

concentrations of methionine, CAATCH1 effectively shuts down ionic fluxes via its channel properties (Feldman *et al.*, 2000, *J. Biol. Chem.* 275(32):24518-24526; Quick, M. and B.R. Stevens, 2001, *J. Biol. Chem.* 276(36):33413-33418).

Compared to conventional organic pesticides, the use of biodegradable small molecules, such as amino acids, as pesticides is highly desirable, owing to the safety of such compounds to humans, animals, and the environment.

Brief Summary

The present invention provides materials and methods for pest control. In a preferred embodiment, the subject invention overcomes drawbacks inherent in the prior art by providing pesticidal compositions that contain one or more compounds that interact with organic solute transporter/ligand-gated ion channel multifunction polypeptides in the pest. Upon exposure, ingestion, or other means of absorption by a target pest, these compositions either compromise pest growth and/or cause the death of the pest. In a preferred embodiment, the compositions of the subject invention contain one or more amino acids and/or amino acid analogs.

In a preferred embodiment, the materials and methods of the subject invention achieve pest control by disrupting the function of a newly discovered class of multifunction solute transporter/ligand-gated ion channel proteins (the CAATCH1 class of proteins). The CAATCH1 proteins of *Aedes aegypti* (yellow fever mosquito), *Leptinotarsa decemlineata* (Colorado potato beetle), and *Manduca sexta* (tomato hornworm) exemplify this class of proteins. In accordance with the subject invention, CAATCH1 has been found to be a key protein in regulating electrolyte and organic solute fluxes, especially in those pests with midguts exhibiting an alkaline pH. The unanticipated biochemical, physiological, and molecular properties of CAATCH1 indicated that it is a multi-function protein that mediates amino acid uptake in a manner that is thermodynamically uncoupled to ion electrochemical potentials, and furthermore that CAATCH1 simultaneously functions as an amino acid-modulated gated alkali cation channel serving at least Na^+ and K^+ . In accordance with the subject invention, effective pest control is achieved by disrupting the function of a CAATCH1 protein and/or related molecules.

As described herein, the function of the CAATCH1 proteins, and related proteins, can be disrupted in a number of ways to achieve the desired pest control. For example, in accordance

with the subject invention, it has been found that small molecules, which interfere with these proteins, can be administered to a pest, or its situs, to achieve pest control. Specifically exemplified herein is the use of the amino acids methionine or leucine (and/or analogs thereof) to control pests. These compounds can be administered in a wide range of ways including the application of compositions comprising these individual amino acids (and/or their analogs, or bound to other molecules, for example by amide bonds), application of polypeptides which are made up of an abundance of these pesticidal amino acids, and providing a transgenic plant which expresses a pesticidal amount of the amino acids as either free amino acids, as salts of amino acids or their analogs, or bound to other molecules, or existing in polypeptides.

In a preferred embodiment, the methods of the subject invention involve delivering to or applying to a pest a composition that comprises methionine or leucine, or an analog thereof. Exemplary analogs include, but are not limited to, methionine esters, leucine esters, D-methionine, D-leucine, D-tert-leucine, L-tert-leucine, DL-methionine, DL-leucine, L-methioninol, L-leucininol, L- or D-methioninemethylsulfoniium chloride, small methionyl peptides, low molecular weight leucyl peptides, alphaketoisocaproic acid, and the like. Likewise, methyl or ethyl esters of such compounds are also contemplated to be useful, as are keto derivatives of such compounds. In a specific embodiment, insect pests are controlled by administering methionine to the insects. Alternatively, other amino acids such as histidine, glycine, threonine, or alanine can also be used. For all embodiments involving compounds with chiral centers, the L-, D-, DL-, or partially racemic forms are also contemplated to be useful. Alternatively, amino acid analogs that do not possess chiral centers are contemplated to be useful.

Additional target crops to be protected within the scope of the present invention comprise, e.g., the following species of plants:

Although it is believed that the administration of amino acids to achieve pest control according to the subject invention is effective as a result of the disruption of particular proteins as described herein, one aspect of the subject invention is simply the control of pests by administering amino acids or these analogs (regardless of the specific mechanism involved).

One aspect of this invention contemplates altering methionine levels in pests by manipulating biochemical pathways leading to methionine production in target pests. Such

manipulations include, but are not limited to, precursors and cofactors of methionine metabolic biosynthesis pathways.

The subject invention provides various other alternative approaches to disrupting the function of the unique newly discovered class of multifunction transporter/channel proteins.

5 These other approaches include, for example, the use of interfering RNA (RNAi), gene silencing techniques, and antisense polynucleotides. In a specific embodiment, the antisense molecules are complementary to contiguous nucleotide sequences of at least about 15 nucleotides from SEQ ID NO:1. These antisense constructs can be used to "down-regulate" the expression of CAATCH1, and/or related proteins, in a particular cell. Alternatively, antisense constructs
10 complementary to a contiguous nucleotide sequence from the CAATCH1 promoter sequence may also be used to regulate the activity of CAATCH1 and/or related proteins. Antisense constructs are well-known in the art and include the use of antisense mRNA to reduce the transcription or translation or otherwise impair the net production of the encoded polypeptide.

The subject invention also provides methods for identifying compounds which regulate, alter, or modulate the biochemical and/or physiological functional activity of a CAATCH1 polypeptide or polynucleotide (or related molecules). In one embodiment this method comprises exposing a cell that expresses a CAATCH1 polypeptide to at least one compound or signal whose ability to modulate the activity of the CAATCH1 polypeptide is sought to be determined, and thereafter monitoring the cell for a change that is a result of the modulation of activity of
15 CAATCH1 or related polypeptide(s). Such an assay is particularly contemplated to be useful in the identification of agonists, antagonists and/or allosteric modulators of CAATCH1.

A further aspect of the invention provides methods for screening compounds (*e.g.*, synthetic peptides, peptide analogs, peptidomimetics, small molecule inhibitors, *etc.*) which inhibit or reduce the binding of a CAATCH1 polypeptide. According to this embodiment,
25 screening for chemical or biochemical entities may be performed *e.g.*, by means of a cell-based assay, an *in vitro* assay for CAATCH1 function and/or rational pesticidal formulation or amino acid transporter-active analogs, drugs, or compounds. Cell-based assays for screening can be designed *e.g.*, by constructing cell lines in which the expression of a reporter protein, *i.e.* an easily assayable protein, is dependent on CAATCH1 activity. Such an assay enables the
30 detection of compounds that directly antagonize CAATCH1, or compounds that inhibit other

cellular functions required for the activity of CAATCH1. Compounds may also be identified which recognize or inhibit amino acid or other solute transport via the CAATCH1 polypeptide or modify ion flux through the CAATCH1 polypeptide. Example ions include, but are not limited to, N^+ , K^+ , H^+ , OH^- , Cl^- , bicarbonate, and carbonate.

5 In another aspect, the present invention provides an antibody that is immunoreactive with a transporter/channel polypeptide of the invention. Reference to antibodies includes whole polyclonal and monoclonal antibodies, and parts thereof, either alone or conjugated with other moieties. The monoclonal antibodies of the present invention can be used in standard immunochemical procedures, such as immunoprecipitation, ELISA and Western blot methods.
10 Also, immunoabsorbent protocols may be used in purifying native or recombinant peptide species or synthetic or natural variants thereof.

Advantageously, the amino acid-based targeted pesticides of the subject invention are environmentally safe due to target selectivity, low toxicity to humans and pets, and their biodegradation by environmentally friendly naturally occurring microorganisms. Also, the use of these pesticides is compatible with the use of natural enemies of pests (e.g., parasitoids and predators). In fact, L-methionine is an "essential or indispensable" amino acid in humans, meaning that L-methionine is not synthesized by the body but instead is a required nutrient in the diet needed to sustain human life (Fuller, M.F., 2000. "Protein and amino acid requirements", pp.287-304 In: *Biochemical and Physiological Aspects of Human Nutrition*, M.H. Stipanuk, ed. W.B. Saunders Co., Philadelphia). Since it is a naturally occurring substance, the use of this compound and related amino acids contribute to a sustainable, pesticide-free food supply, and preserve the environment by reducing the reliance on traditional pesticides.

Brief Description of the Drawings

25 **FIG. 1** shows the efficacy of 10% L-methionine as a *Manduca sexta* pesticide on a defined diet.

FIG. 2 shows the effect of L-methionine concentration as a *Manduca sexta* pesticide as applied to eggplant (*Solanum melongena*) host plants.

30 **FIG. 3** shows the efficacy of L-methionine as a *Manduca sexta* pesticide as applied to eggplant (*Solanum melongena*) host plant leaves using a Silwet-77 solution.

FIG. 4 shows the efficacy of L-methionine as a *Leptinotarsa decemlineata* pesticide as applied to eggplant (*Solanum melongena*) host plants

FIG. 5 shows that L-methionine treatment for *L. decemlineata* or *M. sexta* does not adversely affect host plant fruit yields or weights of host eggplant (*Solanum melongena*) grown in the field.

FIG. 6 shows the efficacy of L-methionine as an *Aedes aegypti* pesticide.

FIG. 7 shows the pesticidal specificity of L-methionine compared to L-proline on *Aedes aegypti* survival.

Brief Description of Sequences

SEQ ID NO: 1 is a DNA sequence, including 5' and 3' untranslated regions and the open reading frame, that encodes the CAATCH1 protein of *Manduca sexta*, *Aedes aegypti*, and *Leptinotarsa decemlineata*.

SEQ ID NO: 2 is the predicted amino acid sequence of the CAATCH1 protein of *Manduca sexta*, *Aedes aegypti*, and *Leptinotarsa decemlineata*, based on the open reading frame of SEQ ID NO:1.

SEQ ID NOS: 3-27 are primers used as described herein to clone CAATCH1 genes.

Detailed Disclosure

The subject invention provides novel pest control compositions, and methods for using such compositions. In one embodiment, the subject compositions can be used to control pests of agricultural crops. These pests, include, for example, coleopterans (beetles), lepidopterans (caterpillars), and mites. The compounds of the subject invention can also be used to control household pests including, but not limited to, ants, termites, and cockroaches. The compounds can also be used to control mosquitoes such as *Aedes aegypti*, *Anopheles gambiae*, and *Culex spp.* Other biting pests such as flies, fleas, ticks, and lice can also be controlled using the compounds and methods of the subject invention. In a preferred embodiment, the materials and methods of the subject invention are used to control pests which have an alkaline gut compartment. Advantageously, the methods and materials of the subject invention provide a novel environmentally friendly, and highly effective approach to controlling pests.

In a preferred embodiment, the methods of the subject invention involve providing access to, or applying to a pest, a composition that comprises methionine or leucine, or an analog thereof. Exemplary analogs include, but are not limited to, methionine esters, leucine esters, D-methionine, D-leucine, D-tert-leucine, L-tert-leucine, DL-methionine, DL-leucine, L-methioninol, L-leucininol, L- or D-methioninemethylsulfoniium chloride, small methionyl peptides, low molecular weight leucyl peptides, alphaketoisocaproic acid, and the like. Likewise, methyl or ethyl esters of such compounds are also contemplated to be useful, as are keto analogs of such compounds. In a specific embodiment, pests are controlled by administering methionine to the pests. Alternatively, other amino acids such as histidine, glycine, threonine, or alanine can also be used. For all embodiments involving compounds with chiral centers, the L-, D-, DL-, or partially racemic forms are also contemplated to be useful. Alternatively, amino acid analogs that do not possess chiral centers are contemplated to be useful.

Methionine levels can be manipulated in the pest via at least two methionine synthetases (Jaffe, J.J. and L.R. Chrin, 1979, *J. Parasitol.* 65:550-554), E.C.2.1.1.13 and E.C.2.1.1.5. A further aspect of this invention concerns approaches to manipulate methionine levels in pest via metabolic pathways. In a one embodiment, pests are controlled by altering the precursors and cofactors of methionine enzymatic biosynthesis, which includes but is not limited to manipulating levels of betaine, homocysteine, S-adenosylmethionine, 5-methyltetrahydrofolate, and colbalamine and its derivatives including vitamin B12.

The materials and methods of the subject invention exert their pesticidal activity by disrupting a new category of ion-associated organic solute transporter proteins (CAATCH1). The CAATCH1 proteins of *Manduca Sexta*, *Aedes aegypti*, or *L. decemlineata* typify this new CAATCH1 class of proteins. In the *M. Sexta* alkaline midgut this protein displays a dual role as an amino acid transporter and an amino acid ligand-gated ion channel. The *M. Sexta* CAATCH1 protein represents the first of its kind for a new class of membrane proteins. In a unique manner, the substrate selectivity of CAATCH1 is modulated by physiological conditions depending on the presence of either K^+ or Na^+ ion at alkaline pH. L-methionine modulates the CAATCH1 reaction mechanism by binding to CAATCH1 and blocking cation current *via* the ion channel aspect of the CAATCH1 polypeptide. By interfering with the combined amino acid

transporting/ligand-gated K^+/Na^+ ion channel properties of CAATCH1, the homeostasis-maintaining role of the protein is perturbed and leads to disruption of fluid, electrolyte and nutrient movements.

Although the disruption of the function of the CAATCH1 protein from *M. sexta*, *A. aegypti*, or *Leptinotarsa decemlineata* is specifically exemplified herein, the materials and methods of the subject invention encompass the analogous disruption of other related transporters, channels, or multifunction transporter/channel proteins in other target pests. Further, the subject invention contemplates direct inhibition of the CAATCH1 class of proteins as well as the disruption of the cascade of cellular activities associated with these unique transporter proteins. In its role as a nutrient transporter, the CAATCH1 protein has been shown to be primarily responsible for proline uptake (Feldman *et al.*, 2000, *supra*), while in the presence of extremely low concentrations of methionine, CAATCH1 effectively shuts down ionic fluxes via its channel aspect (Feldman *et al.*, 2000, *J. Biol. Chem.* 275(32):24518-24526; Quick, M. and B.R. Stevens, 2001, *J. Biol. Chem.* 276(36):33413-33418).

As would be appreciated by one skilled in the art having the benefit of the instant disclosure, pest control can be achieved in a number of ways. For example, compositions comprising individual pesticidal amino acids can be administered directly to the pests. Alternatively, these amino acids may be provided as components of polypeptides, other covalently linked polymers of amino acids or their analogs, or salts of amino acids or their analogs, which are then degraded to the individual amino acids or their analogs in the pest gut and/or prior to entry into the pest gut. These polypeptides or covalently linked compounds may be produced synthetically or recombinantly. In the case of recombinant production, the polypeptides may be produced, for example, in a microbial host followed by isolation of the polypeptide for subsequent administration. The microbial produced polypeptide may also be administered directly to the pest and/or its *situs*.

The polypeptides which contain the pesticidal amino acids of the subject invention may also be expressed in plant cells. The polypeptide may be isolated from the plant cell or, preferably, pest control is achieved by a pest ingesting the plant material. Furthermore pest control can be achieved by avoidance behavior whereby the deleterious effect to the host is reduced by, for example, the pest avoiding consumption of the plant.

One method for controlling pests according to the subject invention provides materials and methods for controlling pests by using double-stranded interfering RNA (RNAi), or RNA-mediated interference (RNAi). The use of antisense compounds is also contemplated.

In view of the use of recombinant hosts according to the subject invention, a further aspect of the subject invention is the polynucleotides that encode the pesticidal polypeptides. These polynucleotide sequences can be readily synthesized by a person skilled in the art, and can be used to transform an appropriate prokaryotic or eukaryotic host to enable the host to express the pesticidal compounds. Hosts of particular interest include bacteria, yeasts, viruses, and plants. For each of these hosts, the DNA sequences may be specifically designed by a person skilled in the art to utilize codons known to be optimally expressed in the particular hosts. Advantageous promoters can also easily be employed in the polynucleotide sequences.

Definitions

As used herein, the term “pesticidal” refers to the ability to interfere with a pest’s life cycle in any way that results in an overall reduction in the pest population. For example, the term pesticidal includes inhibition of a pest from progressing from one form to a more mature form, *e.g.*, transition between various larval instars or transition from larva to pupa or pupa to adult. Further, the term “pesticidal” is intended to encompass anti-pest activity during all phases of a pest’s life cycle; thus, for example, the term includes larvacidal, ovicidal, and adulticidal activity. As used herein, the term “pesticidally effective” is used to indicate an amount or concentration of a pesticidal compound which is sufficient to reduce the number of pests in a geographic locus as compared to a corresponding geographic locus in the absence of the amount or concentration of the pesticidal compound. Pests which can be controlled according to the subject invention are invertebrate animal pests of homes, people, and agriculture. “Pest Control” as used herein includes “pesticidal” activity as well as pest aversion activity which causes a pest to avoid deleterious behavior such as a mosquito biting, or a caterpillar eating an agricultural crop.

The word “transform” is broadly used herein to refer to introduction of an exogenous polynucleotide sequence into a prokaryotic or eukaryotic cell by any means known in the art (including for example, direct transmission of a polynucleotide sequence from a cell or virus

particle, transmission by infective virus particles and transmission by any other nucleotide-bearing construct) resulting in a permanent or temporary alteration of genotype.

The term "upstream" refers to DNA adjacent to the 5' portion of a DNA sequence; whereas, the term "downstream" refers to DNA adjacent to the 3' portion of a DNA sequence.

5 A "transgenic cell" is any cell derived or regenerated from a transformed cell or derived from a transgenic cell. Exemplary transgenic cells include plant calli derived from a transformed plant cell and particular cells such as leaf, root, stem, *e.g.*, somatic cells, or reproductive (germ) cells obtained from a transgenic plant.

10 As used herein, a "transgenic plant" is a plant or progeny thereof derived from a transformed plant cell or protoplast, wherein the plant DNA contains an introduced exogenous DNA molecule not originally present in a native, non-transgenic plant of the same strain.

15 In accordance with the present invention, nucleic acid sequences include and are not limited to, DNA, including and not limited to cDNA and genomic DNA, and genes; RNA, including and not limited to mRNA and tRNA; antisense sequences; and recombinant vectors, including, for example, plasmids, cosmids, phagemids, artificial chromosomes, phage, viruses, baculoviruses, and the like.

20 The term "gene" is used for simplicity to refer to a functional protein-, polypeptide- or peptide-encoding unit. As will be understood by those in the art, this functional term includes genomic sequences, operon sequences and smaller engineered gene segments that express, or may be adapted to express, proteins, polypeptides or peptides.

As used herein with regard to molecular biology, a "vector" is a DNA molecule capable of replication in a host cell and/or to which another DNA segment can be operatively linked so as to bring about replication of the attached segment. A plasmid is an exemplary vector. Note, however, that in entomology terms, a "vector" is a disease transmitting insect.

25 The terms "polypeptide", "peptide", and "protein" as used herein refer to amino acid sequences of two or more amino acids.

The one-letter symbol for the amino acids used herein is well known in the art. For convenience, the relationship of the three-letter abbreviation and the one-letter symbol for amino acids is as follows:

	Ala	A	Leu	L
	Arg	R	Lys	K
	Asn	N	Met	M
	Asp	D	Phe	F
5	Cys	C	Pro	P
	Gln	Q	Ser	S
	Glu	E	Thr	T
	Gly	G	Trp	W
	His	H	Tyr	Y
10	Ile	I	Val	V

CAATCH1

In a preferred embodiment, the subject invention provides materials and methods for achieving pest control by disrupting the function of the new class of transporter/channel proteins exemplified by the *Manduca Sexta* CAATCH1 protein. Figure 1 shows a CAATCH1 polynucleotide. The full length coding sequence of 2858 nucleotides (nt) contains an open reading frame of 1899 nt encoding a predicted 633 amino acid sequence. The nucleotide sequence of this CAATCH1 cDNA open reading frame is about 35-50% identical to a family of clones representing membrane transporters of neurotransmitters and amino acids in a number of species. The predicted amino acid sequence of this CAATCH1 protein is 35-40% identical to most of the stated transporters.

The most closely related clone appears to be the *M. sexta* KAAT1 leucine transporter (Castagna *et al.*, 1998, *Proc. Natl Acad Sci. USA* 95:5395-5400), which is 92% identical overall with the CAATCH1 nucleotide sequence and 90% identical in the overall predicted amino acid sequence. Nonetheless, several regions yield conspicuous differences between CAATCH1 and KAAT1. Notably, the amino acid differences are particularly divergent within or near predicted transmembrane domains #6, #11 and #12, and their adjacent hydrophilic cytosolic C- and N-terminal regions. For example, within residues #496-#577, 25 amino acids (or 31%) diverge often with striking differences.

Although the nucleotide and polypeptide sequences of KAAT1 and CAATCH1 are related, their notable differences are manifested in their striking physiological/functional differences. Thus, the structural and functional differences distinguish them as unique entities.

The DNA sequence information provided herein facilitates the preparation of DNA (or RNA) sequences having the ability to specifically hybridize to nucleic acid sequences encoding portions of the *M. Sexta* CAATCH1 gene, and related genes. The ability of such nucleic acid probes to specifically hybridize to the corresponding *CAATCH1* nucleic acid sequences lend them particular utility in the identification, isolation, and/or characterization of related genes and proteins. Such related proteins may be obtained from *M. sexta*, other members of the genus *Manduca*, *Aedes aegypti* and other mosquitoes, *Leptinotarsa decemlineata*, other insects, or from virtually any other source, plant, animal, fungal, or microbial, from which amino acid transporter/ion channel polypeptides may be isolated that show similarity or homology to the CAATCH1 polypeptide isolated from *M. sexta*, *Aedes aegypti*, or *Leptinotarsa decemlineata*.

In addition to *Manduca* CAATCH1, genes encoding CAATCH1 in the mosquito, *Aedes aegypti* and in the Colorado potato beetle, *Leptinotarsa decemlineata* have been cloned and sequenced and have been found to have the same sequence as the *M. Sexta* CAATCH1 sequence.

Pesticidal Compositions and Methods of Use

The subject invention provides novel pest control compositions, and methods for using such compositions. In an embodiment that is preferred because of its effectiveness, simplicity, and eco-friendliness, a compound, which disrupts the critical functions of transporter proteins, can be administered to a target pest. Specifically exemplified herein is the pesticidal use of leucine and methionine, and/or analogs thereof.

In one embodiment, the subject compositions can be used to control pests of agricultural crops. These pests, include, for example, coleopterans (beetles), lepidopterans (caterpillars), mites, and nematodes. The Coleopterans include numerous beetle species including ground beetles, reticulated beetles, skin and larder beetles, long-horned beetles, leaf beetles, weevils, bark beetles, ladybird beetles, soldier beetles, stag beetles, water scavenger beetles, and a host of other beetles.

Particularly important among the Coleoptera are the agricultural pests included within the infraorders *Chrysomeliformia* and *Cucujiformia*. Members of the infraorder *Chrysomeliformia*, including the leaf beetles (*Chrysomelidae*) and the weevils (*Curculionidae*), are particularly problematic to agriculture, and are responsible for a variety of insect damage to crops and plants.

5 The infraorder *Cucujiformia* includes the families *Coccinellidae*, *Cucujidae*, *Lagridae*, *Meloidae*, *Rhipiphoridae*, and *Tenebrionidae*. Within this infraorder, members of the family *Chrysomelidae* (which includes the genera *Exema*, *Chrysomela*, *Oreina*, *Chrysolina*, *Leptinotarsa*, *Gonioctena*, *Oulema*, *Monozia*, *Ophraella*, *Cerotoma*, *Diabrotica*, and *Lachnaia*), are well-known for their potential to destroy agricultural crops.

10 The compositions of the subject invention can be used as pesticidal formulations against members of the Order Lepidoptera. Likewise, the materials and methods of the subject invention can be used to control mosquitoes (including those in the genera *Aedes*, *Anopheles*, and *Culex*).

Crops which can be protected according to the subject invention as well as pests of these crops include those which are listed in published PCT Application Nos. WO 98/44137 and WO 15 96/10083 which are both incorporated herein by reference.

In a preferred embodiment, the pests controlled according to the subject invention have an alkaline gut compartment. As used herein, reference to alkaline gut compartment means that the typical pH of the gut compartment is greater than 7.0. The alkaline midgut of *M. Sexta* is an example of such an alkaline gut compartment. Other pests having an alkaline gut compartment are set forth in Example 15. Further pest targets according to the subject invention include pests 20 which have a V-type ATPase either expressed as a protein, mRNA, or genomic DNA.

Thus, amino acid-rich compositions, and particularly those that include methionine or leucine (or analogs thereof) in their formulation can be used as pesticides for application to field crops, including but not limited to rice, wheat, alfalfa, corn (maize), soybeans, tobacco, tomato, 25 potato, barley, canola (rapeseed), sugarbeet, sugarcane, flax, rye, oats, cotton, sunflower; grasses, such as pasture and turf grasses; fruits, citrus, nuts, trees, shrubs and vegetables; as well as ornamental plants, cacti, succulents, and the like. Preferred embodiments include plants selected from the group consisting of maize, sorghum, wheat, sunflower, tomato, cole crops, cotton, rice, soybean, sugar beet, sugarcane, tobacco, barley, and oilseed rape. In a particularly preferred 30 embodiment, the plant is a maize plant.

Additional target crops to be protected within the scope of the present invention comprise, e.g., the following species of plants:

Cereals (wheat, barley, rye, oats, rice, sorghum and related crops), beet (sugar beet and fodder beet), forage grasses (orchard grass, fescue, and the like), drupes, pomes and soft fruit (apples, pears, plums, peaches, almonds, cherries, strawberries, raspberries and blackberries), leguminous plants (beans, lentils, peas, soybeans), oil plants (rape, mustard, poppy, olives, sunflowers, coconuts, castor oil plants, cocoa beans, groundnuts), cucumber plants (cucumber, marrows, melons) fiber plants (cotton, flax, hemp, jute), citrus fruit (oranges, lemons, grapefruit, mandarins), vegetables (spinach, lettuce, asparagus, cabbages and other Brassicae, onions, tomatoes, potatoes, paprika), lauraceae (avocados, carrots, cinnamon, camphor), deciduous trees and conifers (e.g. linden-trees, yew-trees, oak-trees, alders, poplars, birch-trees, firs, larches, pines), or plants such as maize, tobacco, nuts, coffee, sugar cane, tea, vines, hops, bananas and natural rubber plants, as well as ornamentals (including composites).

Pesticidal compounds of the subject invention can be used, alone or in combination with other pesticides, to control one or more non-mammalian pests. These pests may be, for example, those listed in Table 1. Activity can readily be confirmed using the bioassays provided herein, adaptations of these bioassays, and/or other bioassays well known to those skilled in the art.

Table 1. Examples of Target pest species

ORDER/Common Name	Latin Name
LEPIDOPTERA	
European Corn Borer	<i>Ostrinia nubilalis</i>
European Corn Borer resistant to Cry1A	<i>Ostrinia nubilalis</i>
Black Cutworm	<i>Agrotis ipsilon</i>
Fall Armyworm	<i>Spodoptera frugiperda</i>
Southwestern Corn Borer	<i>Diatraea grandiosella</i>
Corn Earworm/Bollworm	<i>Helicoverpa zea</i>
Tobacco Budworm	<i>Heliothis virescens</i>
Tobacco Budworm Rs	<i>Heliothis virescens</i>

Table 1. Examples of Target pest species

ORDER/Common Name	Latin Name
Sunflower Head Moth	<i>Homeosoma ellectellum</i>
Banded Sunflower Moth	<i>Cochylis hospes</i>
Argentine Looper	<i>Rachiplusia nu</i>
Cabbage Looper	<i>Trichoplusia ni</i>
Spilosoma	<i>Spilosoma virginica</i>
Bertha Armyworm	<i>Mamestra configurata</i>
Diamondback Moth	<i>Plutella xylostells</i>
COLEOPTERA	
Red Sunflower Seed Weevil	<i>Smicronyx fulvus</i>
Sunflower Stem Weevil	<i>Cylindrocopturus adpersus</i>
Sunflower Beetle	<i>Zygoramma exclamationis</i>
Canola Flea Beetle	<i>Phyllotreta cruciferae</i>
Western Corn Rootworm	<i>Diabrotica virgifera virgifera</i>
DIPTERA	
Hessian Fly	<i>Mayetiola destructor</i>
HOMOPTERA	
Greenbug	<i>Schizaphis graminum</i>
HEMIPTERA	
Lygus Bug	<i>Lygus lineolaris</i>
NEMATODA	
	<i>Heterodera glycines</i>

The pesticidal compounds of the present invention may be provided in a variety of ways. In one example the compounds are provided as a polypeptide, the amino acid sequence of which includes one or more pesticidal amino acids of the present invention. In various embodiments, two or more of the pesticidal amino acids are linked, for example, by peptide bonds between the N-terminus of one portion and the C-terminus of another portion. In other aspects, one or more of the pesticidal polypeptides can be linked to one or more heterologous peptides or proteins to form pesticidal fusion polypeptides. Molecules comprising such portions linked by hydrocarbon

linkages are also provided. Derivatives of the foregoing fusion proteins are also provided (*e.g.*, branched, cyclized, or C-terminal chemically modified, etc.).

Virtually any polypeptide-encoding DNA sequence may be fused to the sequences disclosed herein in order to encode a fusion protein. This includes DNA sequences that encode targeting peptides, proteins for recombinant expression, proteins to which one or more targeting peptides are attached, protein subunits, and the like. Such modifications to primary nucleotide sequences to enhance, target, or optimize expression of the gene sequence in a particular host cell, tissue, or cellular localization, are well known to those of skill in the art of protein engineering and molecular biology. Both N-terminal and C-terminal fusion proteins are contemplated.

Derivation of the pesticidal compounds with long chain hydrocarbons will facilitate passage through the cuticle into the pest body cavity. Therefore, in a further embodiment, the subject invention provides compositions comprising the pesticidal compounds bound to lipids or other carriers.

In addition to the peptide compounds described herein, the subject invention also contemplates that other sterically similar analog compounds may be formulated to mimic the key portions of the peptide structure. Such compounds, which may be termed peptidomimetics, may be used in the same manner as the peptides of the invention and hence are also functional equivalents. The generation of a structural functional equivalent analog may be achieved by the techniques of modeling and chemical design known to those of skill in the art. It will be understood that all such sterically similar analog constructs fall within the scope of the present invention.

The subject invention further contemplates the use of peptide nucleic acids (PNAs) in the practice of the methods of the invention. PNA is a DNA mimic in which the nucleobases are attached to a pseudopeptide backbone. PNA may be utilized in a number of methods that traditionally have used RNA or DNA. Often PNA sequences perform better in techniques than the corresponding RNA or DNA sequences and have utilities that are not inherent to RNA or DNA. A review of PNA including methods of making, characteristics of, and methods of using, is provided by Corey (1997) and is incorporated herein by reference.

Any formulation methods known to those of skill in the art may be employed using methionine or leucine (or other pesticidal compounds as described herein) as an active ingredient. It may be desirable to formulate the amino acid composition alone, or alternatively, by addition of the methionine or leucine composition to existing pesticidal preparations for a combination or synergistic approach to eradicating target pests. For example, methionine or leucine can be added to whole cell preparations, cell extracts, cell suspensions, cell homogenates, cell lysates, cell supernatants, cell filtrates, or cell pellets of cell cultures of pesticide-producing microorganisms. In particular, it may be desired to supplement bacterial cell cultures such as those of *B. thuringiensis* that express one or more δ -endotoxins. The methods for preparing such formulations are known to those of skill in the art, and include, e.g., desiccation, lyophilization, homogenization, extraction, filtration, encapsulation centrifugation, sedimentation, or concentration of one or more cultures of bacterial cells, such as *B. thuringiensis* cells.

In one embodiment, the pesticide composition comprises an oil flowable suspension comprising a methionine or leucine composition. For example, in some embodiments, oil flowable or aqueous solutions may be formulated to contain lysed or unlysed bacterial cells, spores, or crystals which contain one or more pesticidally-active crystal proteins in combination with a methionine or leucine composition. Preferably the cells that contain the crystal protein component of the formulation are *B. thuringiensis* cells; however, any such bacterial host cell expressing one or more crystal-protein encoding polynucleotides is contemplated to be useful, such as *Bacillus* spp., including *B. megaterium*, *B. subtilis*; *B. cereus*, *Escherichia* spp., including *E. coli*, and/or *Pseudomonas* spp., including *P. cepacia*, *P. aeruginosa*, and *P. fluorescens*.

In a further embodiment, the pesticide may be formulated as a water dispersible granule or powder. This granule or powder may comprise one or more amino acids, and particularly, the amino acid methionine or leucine (or analogs thereof).

The pesticide compositions of the present invention may also comprise a wettable powder, spray, emulsion, colloid, aqueous or organic solution, dust, pellet, or colloidal concentrate. Dry forms of the pesticidal compositions may be formulated to dissolve immediately upon wetting, or alternatively, dissolve in a controlled-release, sustained-release, or other time-dependent manner.

Alternatively, the pesticidal composition may comprise an aqueous solution. Such aqueous solutions or suspensions may be provided as a concentrated stock solution which is diluted prior to application, or alternatively, as a diluted solution ready-to-apply. Such compositions may be formulated in a variety of ways. They may be employed as wettable
5 powders, granules or dusts, by mixing with various inert materials, such as inorganic minerals (silicone or silicon derivatives, phyllosilicates, carbonates, sulfates, phosphates, and the like) or botanical materials (powdered corncobs, rice hulls, walnut shells, and the like). The formulations may include spreader-sticker adjuvants, stabilizing agents, other pesticidal additives, or surfactants. Liquid formulations may be employed as foams, suspensions,
10 emulsifiable concentrates, or the like. The ingredients may include rheological agents, surfactants, emulsifiers, dispersants, or polymers.

The compositions may be formulated prior to administration in an appropriate means such as lyophilized, freeze-dried, microencapsulated, desiccated, or in an aqueous carrier, medium or suitable diluent, such as saline or other buffer. Suitable agricultural carriers can be
15 solid or liquid and are well known in the art. The term "agriculturally-acceptable carrier" covers all adjuvants, *e.g.*, inert components, dispersants, surfactants, tackifiers, binders, *etc.* that are ordinarily used in pesticide formulation technology.

The pesticidal compositions of this invention can be applied to the environment of the target pest, typically onto the foliage of the plant or crop to be protected, by conventional
20 methods such as spraying. Other application techniques, *e.g.*, dusting, sprinkling, soaking, soil injection, soil tilling, seed coating, seedling coating, spraying, aerating, misting, atomizing, and the like, are also feasible and may be required under certain circumstances such as *e.g.*, pests that cause root or stalk infestation, or for application to delicate vegetation or ornamental plants. These application procedures are also well known to those of skill in the art.

25 To control mosquito larvae, the compositions may be applied to standing water. In this context, the composition may be, for example, a yeast or algae (or other mosquito or mosquito larvae food) which has been mixed with, or transformed to express, a pesticidal compound of the subject invention.

In the case of termites, the pesticidal compounds (such as methimine) can be used to form a barrier or, preferably are incorporated into a food source such as a wood article. The food source could be used, for example, as the toxicant in a Sentricon™ system.

The pesticidal compositions of the invention may be employed in the method of the invention singly or in combination with other compounds, including, but not limited to, other pesticides.

Regardless of the method of application, the amount of the active component(s) are applied at a pesticidally-effective amount, which will vary depending on factors such as, for example, the specific pests to be controlled, the specific plant or crop to be treated, the environmental conditions, and the method, rate, and quantity of application of the pesticidally-active composition.

The concentration of pesticidal composition that is used for environmental, systemic, or foliar application will vary widely depending upon the nature of the particular formulation, means of application, environmental conditions, and degree of biocidal activity. Typically, the composition will be present in the applied formulation at a concentration that provides the amino acid composition to the pest in a range of from about 0.1% by weight up to and including about 99% by weight. Formulations of the compositions may be from about 0.01% to about 1% by weight, or from 1% to about 75% or more by weight, with some formulations generally comprising from about 5% to about 50% or more of the active ingredient by weight. Formulations which comprise intact bacterial cells will generally contain from about 10^4 to about 10^{12} cells/mg of the final formulation. The pesticidal formulation may be administered to a particular plant or target area in one or more applications as needed, with a typical field application rate per hectare ranging on the order of from about 1 g to about 5 kg, or more of active ingredient.

Transformed Host Cells and Transgenic Plants

A preferred embodiment of the subject invention provides transformed host cells and transgenic plants that express amino acids at levels that are pesticidal to pests feeding on such cells or plants. In a specific embodiment, methionine-encoding polynucleotides are introduced into cells to increase, alter, or affect the overall concentration of methionine within the cells.

The recombinant host may be, for example, prokaryotic or eukaryotic cells such as yeast or

algae. The transformed hosts can be applied to pest habitats, such as bodies of water inhabited by mosquito larvae. Ingestion of the transformed host by a pest species would lead to control of the pest by the pesticidal polypeptide.

Technology for introduction of polynucleotides into cells is well known to those of skill in the art. Four general methods for delivering a gene into cells have been described: (1) chemical methods; (2) physical methods such as microinjection, electroporation and the gene gun; (3) viral vectors; and (4) receptor-mediated mechanisms. Methods for DNA transformation of plant cells include *Agrobacterium*-mediated plant transformation, protoplast transformation, gene transfer into pollen, injection into reproductive organs, injection into immature embryos and particle bombardment.

Bacteria, yeasts, algae, plants, and viruses each may be used in the production of pesticidal polypeptides for further use, or these hosts can be used as vehicles for direct application of pesticidal polypeptides to the target pest. Plants can be transformed to render them toxic to a target pest species that feeds on the transformed plant. In this way, the plant may also be rendered undesirable as a food source thus reducing damage to the plant.

Typically, a gene of interest is introduced between the transcriptional and translational initiation region and the transcriptional and translational termination region, so as to be under the regulatory control of the initiation region. This construct is included in a plasmid, which includes at least one replication system. Where integration is desired, the plasmid will desirably include a sequence homologous with the host genome.

The transformants can be isolated in accordance with conventional ways, usually employing a selection technique, which allows for selection of the desired organism as against unmodified organisms or transferring organisms, when present. The transformants then can be tested for pesticidal activity.

Hosts of particular interest are the prokaryotes and the lower eukaryotes, such as fungi. Illustrative prokaryotes, both Gram-negative and -positive, include *Enterobacteriaceae*; *Bacillaceae*; *Rhizobiaceae*; *Spirillaceae*; *Lactobacillaceae*; and phylloplane organisms such as members of the *Pseudomonadaceae*. Particularly preferred host cells include *Pseudomonas aeruginosa*, *Pseudomonas fluorescens*, *Bacillus thuringiensis*, *Escherichia coli*, *Bacillus subtilis*, and the like.

Among eukaryotes are fungi, such as *Phycomycetes* and *Ascomycetes*, which includes yeast, such as *Schizosaccharomyces*; and *Basidiomycetes*, *Rhodotorula*, *Aureobasidium*, *Sporobolomyces*, *Saccharomyces* spp., and *Sporobolomyces* spp.

A large number of cloning vectors are available for the insertion of foreign genes into many organisms, including plants. In some instances, it may be desirable to provide for regulative expression of the gene encoding the gene of interest where expression is regulated by certain cellular stimuli, environmental conditions, cell cycle, or other inductive or repressive factor(s). This can be achieved, for example, through the incorporation of one or more operators, enhancers, or a region of DNA that binds to an activator or an enhancer, into the genetic element that comprises the gene, so that the genetic construct may be expressed under certain controlled conditions.

Various manipulations may be employed for enhancing the expression of the messenger RNA, particularly by using an active promoter, as well as by employing sequences, which enhance the stability of the messenger RNA. The transcriptional and translational termination region will involve stop codon(s), a terminator region, and optionally, a polyadenylation signal. A hydrophobic "leader" sequence may be employed at the amino terminus of the translated polypeptide sequence in order to promote secretion of the protein across the inner membrane.

A large number of transcriptional regulatory regions are available from a wide variety of microorganism hosts, such as bacteria, bacteriophage, cyanobacteria, algae, fungi, and the like. Various transcriptional regulatory regions include the regions associated with the *trp* gene, *lac* gene, *gal* gene, the λ_L and λ_R promoters, the *tac* promoter, the naturally-occurring promoters associated with the transport protein-encoding gene, where functional in the host. See for example, U. S. Patent Nos. 4,332,898; 4,342,832; and 4,356,270 (each of which is specifically incorporated herein by reference).

Where stable episomal maintenance or integration is desired, a plasmid will be employed which has a replication system that is functional in the host. A large number of plasmids are available, such as pBR322, pACYC184, RSF1010, pR01614, and the like. See for example, U. S. Patent Nos. 4,356,270; 4,362,817; 4,371,625, and 5,441,884, each incorporated specifically herein by reference.

In other embodiments, it is contemplated that certain advantages will be gained by positioning the coding DNA segment under the control of a recombinant, or heterologous, promoter. The use of promoter and cell type combinations for protein expression is generally known to those of skill in the art of molecular biology, for example, see Maniatis *et al.*, 1989.

5 The promoters employed may be constitutive, or inducible, and can be used under the appropriate conditions to direct high level expression of the introduced DNA segment, such as is advantageous in the large-scale production of recombinant proteins or peptides. Appropriate promoter systems contemplated for use in high-level expression include, but are not limited to, the *Pichia* expression vector system (Pharmacia LKB Biotechnology).

10

Following are examples that illustrate procedures for practicing the invention. These examples should not be construed as limiting. All percentages are by weight and all solvent mixture proportions are by volume unless otherwise noted.

15 Example 1- Cloning and Characterization of a Ligand-Gated Ion Channel/Amino Acid Transporter.

A full-length clone was isolated from a midgut cDNA library (in LambdaZap) derived from the fifth instar larval stage of *M. sexta*. The clone included 5' and 3' untranslated regions, in addition to a functional open reading frame. A strategy was employed using inosine degenerate primer touchdown PCR™ combined with specific primer based PCR™ screening of phage lysates to isolate the clone (denoted "CAATCH1"), with subsequent expression and extensive *in vitro* characterization of the expressed polypeptide product in membranes of *Xenopus* oocytes.

20 An initial set of inosine ("I")-containing degenerate primers was designed (Feldman *et al.* JBC, 2000) to target conserved peptide motifs from invertebrate and vertebrate members of a subfamily of Na/Cl dependent transporters serving various neurotransmitters and amino acids (Griffith, J.K. and C.E. Sansom, 1998; <http://www.biology.ucsd.edu/~msaier/transport/classf.html>). All primers are shown in the 5' to 3' direction. The sense primer "S34" (GGIAA(C/T)GTITGG(A/C)G(A/G/C/T)TT(C/T)CC) 25 (SEQ ID NO:3) was based on a GNVWRFP (SEQ ID NO:4) peptide motif, while the antisense 30

primer "S21" (IGC(A/G/T)ATIGCITC(A/G/C/T)GG(A/G)TA) (SEQ ID NO:5) was based on a YP(D/E)AIA (SEQ ID NO:6) peptide motif. Another sense primer "S22" (GGIAA(C/T)GTITGG(G/T)G(A/G/C/T)TT(C/T)CC) (SEQ ID NO:7), a tolerated alternative to S34, was also used in conjunction with S21 antisense primer for initial screening. Another primer set was designed to specifically exclude KAAT1 and other potentially related sequences, including, while amplifying a unique 328 bp segment. In this case, sense primer "S25" (AACACTTGCTGCATCAGTCAAC) (SEQ ID NO:8) and antisense primer "S26" (CTCAAGGAGTTTCGCCCATG) (SEQ ID NO:9). The S25/S26 set was used for subsequent library phage lysate PCRTM screening steps, and was used with the cloned 943 bp fragment to create a 328 bp digoxigenin (DIG)-labeled (Boehringer-Mannheim) dsDNA plaque hybridization probe for the isolation of a single clone. The sequence of the full length clone (SEQ ID NO:1) was determined, including the open reading frame plus the 3' and 5' untranslated regions.

Example 2 - Cloning and Characterization of a Ligand-Gated Ion Channel/Amino Acid Transporter.

A CAATCH1 gene was also cloned from *Aedes aegypti* mosquito larvae mRNA and from *Leptinotarsa decemlineata* Colorado potato beetle larvae mRNA. In each case, the RNA was isolated from excised larval guts or intact larvae. First strand cDNA was produced using reverse transcriptase with either primer "AP" (Gibco Life Technologies) (GGCCACGCGTCGACTAGTACTTTTTTTTTTTTTTTTTT) (SEQ ID NO:10), or PA142_AP"(GACTTCAGGCTAGCATCGATCCATGGGTCGACTTTTTTTTTTTTTTTTTT) (SEQ ID NO:11). CAATCH1-specific fragments were subsequently amplified from the first strand cDNA by PCRTM using a proofreading DNA polymerase mixed with Taq (Invitrogen; Platinum HighFidelity), employing various primer combinations of sense and antisense sequences.

Sense sequences that were used to clone and/or sequence CAATCH1 from *Aedes aegypti* or *Leptinotarsa decemlineata* included the following (all shown in the 5' to 3' direction):

"PA140_AUAP" (GACTTCAGGCTAGCATCGATCCA) (SEQ ID NO:12);

"PA141" (CATCGATCCATGGGTCGAC) (SEQ ID NO:13);

"AUAP (Gibco)" (GGCCACGCGTCGACTAGTAC) (SEQ ID NO:14);

“left1_Forward” (GGCACGAGGTTACTTGTGG) (SEQ ID NO:15);
 “left2_Forward” (CGAGGTTACTTGTGGAGGAA) (SEQ ID NO:16);
 “ATG1_Forward” (TTGTGGACAAAATGAATGACG) (SEQ ID NO:17);
 “ATG2_Forward” (AAAATGAATGACGGCCAAGT) (SEQ ID NO:18);
 5 “S25TrueFOR” (AACACTTGCTGCATCAGTCAAC) (SEQ ID NO:8);
 “uniqueFOR” (TGCTGCATCAGTCAACAACA) (SEQ ID NO:19);
 “TaqFOR” (TCATCCTGCCCCGGTGCTA) (SEQ ID NO:20);
 “1445_FOR” (ACACCGGGTGGACAATATATTCTT) (SEQ ID NO:21);
 “StopORF_FOR” (GGTGCTTACAGGCGTAATATTAATTAG) (SEQ ID NO:22);
 10 “3'UTR_FOR” (CGGTGTCGATATTATACATGTTACGA) (SEQ ID NO:23)
 and “S34” (GGIAA(C/T)GTITGG(A/C)G(A/G/C/T)TT(C/T)CC) (SEQ ID NO:3).

Antisense sequences used to clone and/or sequence CAATCH1 from *Aedes aegypti* or *Leptinotarsa decemlineata* included (all shown from 5' to 3' direction):

“CAATCH1_unique_REVERSE” (AGGAGTTTCGCCCCATTGAG) (SEQ ID NO:24);
 15 “Taq_CAATCH1_REVERSE” (CAAGGAGTTTCGCCCCATTGA) (SEQ ID NO:25);
 “S26_REVERSE” (CTCAAGGAGTTTCGCCCCATTG) (SEQ ID NO:9);
 “CAATCH1_2642_REV” (CTATAAAATCAAACGCAAGACCAT) (SEQ ID NO:26);
 and CAATCH1_2727_REV” (TGTTGCACAGAGTTCTTGAAGATTT) (SEQ ID NO:27).

20 Example 3 - DNA Sequence Analysis and Protein Functional Behavior

The complete sequence of the CAATCH1 cDNA clone, including 5' and 3' UTRs and the open reading frame encoding the predicted 633 amino acid polypeptide expression product, were obtained for *Manduca sexta* using a midgut cDNA library in LambdaZap phage. The *Manduca* CAATCH1 Genbank accession No. is AF013963 (SEQ ID NO:1). The cDNA sequences of
 25 CAATCH1 from *Aedes aegypti* larvae and from *Leptinotarsa decemlineata* Colorado potato beetle larvae were each cloned from mRNA, and were determined to be the same as SEQ ID NO:1 and SEQ ID NO:2. Southern blot analysis of *Aedes aegypti* and *Leptinotarsa decemlineata* DNA products from PCRTM reactions obtained using various primers (SEQ ID NOS:3—27) indicated the existence of splice variants in the untranslated regions of CAATCH1
 30 from these organisms. A database analysis of the *Anopheles gambiae* genome project revealed

contiguous portions of CAATCH1 polynucleotides, and furthermore, subsequent analysis of PCRTM products obtained from an *Anopheles gambiae* cDNA library using CAATCH1-specific primers (SEQ NOS:3—27) yielded polynucleotides of the correct predicted size.

The CAATCH1 nucleotide coding sequence within the full length clone of 2858 nt (SEQ ID NO:1) contained an open reading frame encoding a predicted unique polypeptide sequence of 633 amino acids (SEQ ID NO:2). The predicted amino acid sequence of CAATCH1 is compared to related members of a subfamily of membrane proteins that includes Na/Cl activated transporters of neurotransmitters (Quick, M. and B.R. Stevens, 2001, *J. Biol. Chem.* 276(36):33413-33418; Feldman *et al*, 2000, *J. Biol. Chem.* 275(32):24518-24526; Griffith, J.K. and C.E. Sansom, 1998, In: *The Transporter Facts Book*, Academic Press, San Diego, pp. 500) and the *Manduca* KAAAT1 leucine transporter (Castagna *et al.*, 1998, *Proc. Natl. Acad. Sci. USA* 95:5395-5400). Based on hydropathy analysis and established membrane protein structure paradigms (Kyte and Doolittle, 1982), the predicted CAATCH1 protein topology includes 12 putative transmembrane domains, with N- and C- terminal segments residing within the cytosol. Several consensus phosphorylation sites are found within these cytoplasmic segments, and N-linked glycosylation sites exist on the putative extracellular loop between the 3rd and 4th membrane spanning segments. The cytosolic N- and C- terminal regions are relatively rich in proline, acidic, and basic amino residues.

Membrane transporters and ion channels in general can be subcategorized based on thermodynamic properties, substrate selectivities, and reaction mechanism (Gerencser, G.A. and B.R. Stevens, 1994, *J. Exper. Biol.* 196:59-75; Hille, B., 2001, In: *Ion Channels of Excitable Membranes*, 3rd Edition, Sinauer Associates, Inc., Sunderland, Mass, pp 814; <http://www.biology.ucsd.edu/~msaier/transport/>). CAATCH1 cloned from *Manduca sexta* collectively displayed a unique set of properties (Quick, M. and B.R. Stevens, 2001, *supra*; Feldman *et al.*, 2000, *supra*) that have not been described previously for a given related transporter, including at least the following attributes. These same attributes are also assigned to both *Aedes aegypti* and *Leptinotarsa decemlineata* by inference of their sequences being identical to SEQ NO:1. The attributes include: (a) the ability to switch particular amino acid substrate selectivities depending on the activator cation Na⁺ or K⁺, (b) a unique selectivity profile of amino acid-evoked electrical currents, (c) different amino acid substrates directly binding the

protein and differentially affecting the conformational states of CAATCH1, apparent lack of chloride ion as a co-activator, (d) Nernstian cation channel behavior independent of amino acid transporter activity, (e) amino acid-modulated ion channel behavior (especially L-methionine binding to CAATCH1 in the presence of Na^+ which perturbs the charge-voltage relation with a high affinity binding constant, affecting transient currents due to CAATCH1-associated charge transfer across the membrane dielectric field), (f) inhibition of current fluxes as the result of binding of methionine to the protein, (g) thermodynamically uncoupled amino acid transport and ion channel behavior, and (h) all these functions behave optimally at an alkaline pH. CAATCH1 defines a new transport system (Quick, M. and B.R. Stevens, 2001, *supra*; Feldman *et al.*, 2000, *supra*; Castagna *et al.*, 1998, *supra*; Christensen, 1990; Griffith and Sansom, 1998; Kilberg *et al.*, 1993, *Annu. Rev. Nutr.* 13:137-165; Mailliard *et al.*, 1995, *Gastroenterology* 108:888-910; Malandro and Kilberg, 1996; Stevens, B.R., 1992, "Amino Acid Transport in Intestine" *In*: Mammalian Amino Acid Transport: Mechanisms and Control, pp 149-164, M.S. Killberg and D. Haussinger, eds., Plenum, New York). The transport of amino acids by CAATCH1 serves the simultaneous and independent roles of nutrient transporter and amino acid-gated ion channel in *M. sexta*, *A. aegypti*, or *L. decemlineata*.

Example 4 - Identifying CAATCH1 Proteins and Genes.

Homologous polynucleotides and polypeptides can be identified and obtained through several means. The specific genes, or portions thereof, may be constructed synthetically. Variations of these genes may be readily constructed using standard techniques for making point mutations. Also, fragments of these genes can be made using commercially available exonucleases or endonucleases according to standard procedures. For example, enzymes such as *Bal31* or site-directed mutagenesis can be used to systematically cut off nucleotides from the ends of these genes. Also, genes that code for active fragments may be obtained using a variety of other restriction enzymes. Proteases may be used to directly obtain active fragments of these amino acid transport proteins.

Equivalent amino acid transporter/ion channel proteins and/or genes encoding these equivalent amino acid transport/ion channel proteins can also be isolated from, or identified in, other insects, or other species of mosquitoes, hornworms, or caterpillars and/or pest-specific

DNA or RNA libraries. For example, antibodies to the amino acid transport proteins disclosed and claimed herein can be used to identify and isolate other amino acid transport proteins. Specifically, antibodies may be raised to the portions of the amino acid transport proteins that are most constant and most distinct from other proteins. These antibodies can then be used to specifically identify equivalent polypeptides with the characteristic transport activity by immunoprecipitation, enzyme linked immunoassay (ELISA), or Western blotting.

A further method for identifying the amino acid transporter/ion channel proteins and genes of the subject invention is through the use of oligonucleotide probes. These probes are nucleotide sequences having a detectable label. As is well known in the art, if the probe molecule and nucleic acid sample hybridize by forming a strong bond between the two molecules, it can be reasonably assumed that the probe and sample share significant homology. The probe's detectable label provides a means for determining in a known manner whether hybridization has occurred. Such a probe analysis provides a rapid method for identifying, isolating, and/or characterizing amino acid transport protein-encoding genes of the subject invention.

The nucleic acid sequences provided herein have utility as probes or primers in nucleic acid hybridization embodiments. Such sequences may involve the purine and pyrimidine nucleotides and/or inosine substitutions, and may be arranged to encode amino acids according to sequences representing degeneracy of codon usage (Table 2). As such, it is contemplated that nucleic acid segments that comprise a sequence region that consists of at least about 15 nucleotide contiguous sequence that has the same sequence as, or is complementary to, a 15 nucleotide contiguous DNA segment of SEQ ID NO:1 will find particular utility. Longer contiguous identical or complementary sequences, *e.g.*, those of about 20, 50, 100, 500, 1000, 5000, 10,000 *etc.* (including all intermediate lengths and up to and including full-length sequences) will also be of use in certain embodiments. The preferred nucleic acid sequence employed for hybridization studies or assays includes sequences that have, or are complementary to, at least an about 15 to about 29 nucleotide stretch of the sequence, although sequences of about 30 to about 50 nucleotides are also useful. In particular, such polynucleotides preferably comprise contiguous regions from SEQ ID NO:1, and hybridize to the sequence of SEQ ID NO:1 under relatively high stringency hybridization conditions. A variety of hybridization techniques

and systems are known that can be used in connection with the hybridization aspects of the invention, including diagnostic assays such as those described in U. S. Patent No. 4,358,535, incorporated herein by reference.

Table 2								
Amino Acids	Codons							
Alanine	Ala	A	GCA	GCC	GCG	GCU		
Cysteine	Cys	C	UGC	UGU				
Aspartic acid	Asp	D	GAC	GAU				
Glutamic acid	Glu	E	GAA	GAG				
Phenylalanine	Phe	F	UUC	UUU				
Glycine	Gly	G	GGA	GGC	GGG	GGU		
Histidine	His	H	CAC	CAU				
Isoleucine	Ile	I	AUA	AUC	AUU			
Lysine	Lys	K	AAA	AAG				
Leucine	Leu	L	UUA	UUG	CUA	CUC	CUG	CUU
Methionine	Met	M	AUG					
Asparagine	Asn	N	AAC	AAU				
Proline	Pro	P	CCA	CCC	CCG	CCU		
Glutamine	Gln	Q	CAA	CAG				
Arginine	Arg	R	AGA	AGG	CGA	CGC	CGG	CGU
Serine	Ser	S	AGC	AGU	UCA	UCC	UCG	UCU
Threonine	Thr	T	ACA	ACC	ACG	ACU		
Valine	Val	V	GUA	GUC	GUG	GUU		
Tryptophan	trp	W	UGG					
Tyrosine	Tyr	Y	UAC	UAU				

Various degrees of stringency of hybridization can be employed. The more severe the conditions, the greater the complementarity that is required for duplex formation. Temperature, probe concentration, probe length, ionic strength, time, and the like can control severity of conditions. Preferably, hybridization is conducted under moderate to high stringency conditions by techniques well known in the art, as described, for example, Keller, G.H., M.M. Manak, 1987, In: *DNA Probes*, Stockton Press, New York, NY., pp. 169-170.

Examples of various stringency conditions are provided herein. Hybridization of immobilized DNA on Southern blots with ³²P-labeled gene-specific probes can be performed by standard methods (Sambrook *et al.*, 1982, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, New York.). In general, hybridization and subsequent washes can be

carried out under moderate to high stringency conditions that allow for detection of target sequences with homology to the exemplified polynucleotide sequence. For double-stranded DNA gene probes, hybridization can be carried out overnight at 20-25° C below the melting temperature (T_m) of the DNA hybrid in 6X SSPE, 5X Denhardt's solution, 0.1% SDS, 0.1 mg/ml denatured DNA. The melting temperature is described by the following formula (Beltz *et al.*, 1983, *Methods of Enzymology*, 100:266-285, R. Wu, L. Grossman and K. Moldave, eds., Academic Press, New York).

$T_m = 81.5^\circ \text{C} + 16.6 \log[\text{Na}^+] + 0.41(\% \text{G+C}) - 0.61(\% \text{formamide}) - 600/\text{length of duplex in base pairs.}$

Washes are typically carried out as follows:

- (1) twice at room temperature for 15 minutes in 1X SSPE, 0.1% SDS (low stringency wash);
- (2) once at (T_m)-20°C for 15 minutes in 0.2X SSPE, 0.1% SDS (moderate stringency wash).

For oligonucleotide probes, hybridization can be carried out overnight at 10-20°C below the melting temperature (T_m) of the hybrid in 6X SSPE, 5X Denhardt's solution, 0.1% SDS, 0.1 mg/ml denatured DNA. T_m for oligonucleotide probes can be determined by the following formula:

$T_m (^\circ \text{C}) = 2(\text{number T/A base pairs}) + 4(\text{number G/C base pairs})$ (Suggs *et al.*, 1981, *ICN-UCLA Symp. Dev. Biol. Using Purified Genes*, D.D. Brown, ed., Academic Press, New York, 23:683-693).

Washes can be carried out as follows:

- (1) twice at room temperature for 15 minutes 1X SSPE, 0.1% SDS (low stringency wash);
- (2) once at the hybridization temperature for 15 minutes in 1X SSPE, 0.1% SDS (moderate stringency wash).

In general, salt and/or temperature can be altered to change stringency. With a labeled DNA fragment >70 or so bases in length, the following conditions can be used:

Low:	1 or 2X SSPE, room temperature
Low:	1 or 2X SSPE, 42°C

Moderate: 0.2X or 1X SSPE, 65°C

High: 0.1X SSPE, 65°C.

Duplex formation and stability depend on substantial complementarity between the two strands of a hybrid and, as noted above, a certain degree of mismatch can be tolerated.

5 The ability of such nucleic acid probes to specifically hybridize to CAATCH1 protein-encoding sequences enable them to be of use in detecting the presence of complementary sequences in a given sample.

Small nucleic acid segments or fragments may be readily prepared by, for example, directly synthesizing the fragment by chemical means, as is commonly practiced using an automated oligonucleotide synthesizer. Also, fragments may be obtained by application of
10 nucleic acid reproduction technology, such as the PCR™ technology of U. S. Patent Nos. 4,683,195 and 4,683,202 (each incorporated herein by reference), by introducing selected sequences into recombinant vectors for recombinant production, and by other recombinant DNA techniques generally known to those of skill in the art of molecular biology.

Accordingly, the nucleotide sequences of the invention may be used for their ability to selectively form duplex molecules with complementary stretches of DNA fragments. Depending on the application envisioned, one will desire to employ varying conditions of hybridization to achieve varying degrees of selectivity of probe towards target sequence. For applications requiring high selectivity, one will typically desire to employ relatively high stringency
15 conditions to form the hybrids, *e.g.*, one will select relatively low salt and/or high temperature conditions, such as provided by about 0.02 M to about 0.15 M NaCl at temperatures of about 50°C to about 70°C. Such selective conditions tolerate little, if any, mismatch between the probe and the template or target strand, and would be particularly suitable for isolating DNA segments encoding amino acid transport/ion channel proteins. Detection of DNA segments *via*
20 hybridization is well known to those of skill in the art, and the teachings of U. S. Patent Nos. 4,965,188 and 5,176,995 (each incorporated herein by reference) are exemplary of the methods of hybridization analyses.

For some applications, for example, where one desires to prepare mutants employing a mutant primer strand hybridized to an underlying template or where one seeks to isolate amino
30 acid transporter/ion channel protein-encoding sequences from related species, functional

equivalents, or the like, less stringent hybridization conditions will typically be needed in order to allow formation of the heteroduplex. In these circumstances, one may desire to employ conditions such as about 0.15 M to about 0.9 M salt, at temperatures ranging from about 20°C to about 55°C. In any case, it is generally appreciated that conditions can be rendered more stringent by the addition of increasing amounts of formamide, which serves to destabilize the hybrid duplex in the same manner as increased temperature.

In illustrative embodiments, the polynucleotides of the present invention may comprise a nucleic acid sequence having at least about 60% preferably more than about 70%, more preferably more than about 85%, even more preferably more than about 90%, most preferably more than about 95%, even up to and including about 96%, about 97%, about 98%, or about 99% or greater sequence identity with a contiguous nucleic acid sequence of at least about 15 or so nucleotides from SEQ ID NO:1. Of course, the percent identity to a contiguous nucleic acid sequence from SEQ ID NO:1 need not be limited to the specific percentages given, but is also meant to include all integers between about 60% and about 99% identity with a contiguous nucleic acid sequence of at least about 15 nucleotides from SEQ ID NO:1. In fact, all such sequences are contemplated to fall within the scope of the present invention, so long as the particular sequence retains the relevant function.

The CAATCH1 nucleotides and sequences of the present invention include those comprising the stated open reading frame, as well as the 3' and 5' untranslated regions of both the sense and antisense sequences. Furthermore, the polynucleotides of the present invention include all splice variants arising from one or more genomic DNA sequences. Furthermore, both the sense and antisense versions of the primer sequences used to clone or sequence CAATCH1 nucleotide sequences are included in this invention.

Example 5 - Methods for Making and Using Antibodies.

Antibodies, both polyclonal and monoclonal, specific for CAATCH1 and CAATCH1-derived peptides and/or epitopes may be prepared using conventional immunization techniques. A composition containing antigenic CAATCH1 epitopes can be used to immunize one or more animals, such as a rabbit or mouse, which will then produce specific antibodies against epitope-containing CAATCH1 peptides. Polyclonal antisera may be obtained, after allowing time for

antibody generation, simply by bleeding the animal and preparing serum samples from the whole blood.

One feature provided by the present invention is a polyclonal sera that is relatively homogenous with respect to the specificity of the antibodies therein. Typically, polyclonal antisera is derived from a variety of different "clones," *i.e.* B-cells of different lineage. Monoclonal antibodies, by contrast, are defined as coming from antibody-producing cells with a common B-cell ancestor, hence their "mono" clonality. mAbs may be readily prepared through use of well-known techniques, such as those exemplified in U. S. Patent No. 4,196,265, incorporated herein by reference.

In general, both poly- and monoclonal antibodies against these peptides may be used in a variety of embodiments. For example, they may be employed in antibody cloning protocols to obtain cDNAs or genes encoding the peptides disclosed herein or related proteins. They may also be used to inhibit the effects of CAATCH1 in cells or animals.

Example 6 - Epitopic Core Sequences.

The present invention also provides CAATCH1 polypeptide compositions, free from total cells and other polypeptides, which comprise a purified CAATCH1 polypeptide which incorporates an epitope that is immunologically cross-reactive with one or more of the CAATCH1-specific antibodies of the present invention.

As used herein, the term "incorporating an epitope(s) that is immunologically cross-reactive with one or more anti- CAATCH1 antibodies" refers to an antigen which includes a primary, secondary or tertiary structure similar to an epitope located within a CAATCH1 polypeptide. The level of similarity will generally be to such a degree that monoclonal or polyclonal antibodies directed against the CAATCH1 polypeptide will also bind to, react with, or otherwise recognize, the cross-reactive peptide or protein antigen. Various immunoassay methods may be employed in conjunction with such antibodies, such as, for example, Western blotting, ELISA, RIA, and the like, all of which are known to those of skill in the art.

One may employ the methods of Hopp, as taught in U. S. Patent No. 4,554,101, incorporated herein by reference, which teaches the identification and preparation of epitopes from amino acid sequences on the basis of hydrophilicity. The methods described in several

other papers, and software programs based thereon, can also be used to identify epitopic core sequences. The amino acid sequence of these "epitopic core sequences" may then be readily incorporated into peptides, either through the application of peptide synthesis or recombinant technology.

5 As an exemplary embodiment, to conduct a competition study between CAATCH1 and any test compound, one would first label CAATCH1 with a detectable label to enable subsequent identification. One would then incubate the labeled antigen with the other, test, antigen and, after mixing, one would then add the mixture to a known antibody. The ability of the mixture to bind to the antibody would be determined by detecting the presence of the specifically bound
10 label. This value would then be compared to a control value in which no potentially competing (test) antigen was included in the incubation.

The assay may be any one of a range of immunological assays based upon hybridization, and the reactive antigens would be detected by means of detecting their label, *e.g.*, using streptavidin in the case of biotinylated antigens or by using a chromogenic substrate in
15 connection with an enzymatic label or by simply detecting a radioactive or fluorescent label.

A significant reduction in labeled antigen reactivity in the presence of a test antigen is indicative of a test antigen that is "cross-reactive", *i.e.* that has binding affinity for the same antibody. "A significant reduction", in terms of the present application, may be defined as a reproducible (*i.e.* consistently observed) reduction in binding.

Example 7 - Mutagenesis of CAATCH1 Polypeptides and Polynucleotides.

20 In certain embodiments, it is desirable to prepare mutant polypeptides and/or polynucleotides that encode them. Once the structure of the desired peptide to be mutagenized has been analyzed, it may often be desirable to introduce one or more mutations into either the polypeptide sequence, alternatively, into the DNA sequence encoding the CAATCH1-derived polypeptide for the purpose of producing a mutated peptide with altered biological properties.

25 To that end, the present invention encompasses both site-specific mutagenesis methods and random mutagenesis of a nucleic acid segment encoding a transport polypeptide of the present invention. Mutagenesis may be performed in accordance with any of the techniques

known in the art such as and not limited to synthesizing an oligonucleotide having one or more mutations within the sequence of a particular polypeptide.

For example, certain amino acids may be substituted for other amino acids in a protein structure without appreciable loss of interactive binding capacity with structures such as, for example, antigen-binding regions of antibodies or binding sites on substrate molecules.

In making such changes, the hydropathic index of amino acids may be considered. The importance of the hydropathic amino acid index in conferring interactive biologic function on a protein is generally understood in the art. It is accepted that the relative hydropathic character of the amino acid contributes to the secondary structure of the resultant protein, which in turn defines the interaction of the protein with other molecules, for example, enzymes, substrates, receptors, DNA, antibodies, antigens, and the like. Each amino acid has been assigned a hydropathic index on the basis of their hydrophobicity and charge characteristics (Kyte and Doolittle, 1982), these are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (−0.4); threonine (−0.7); serine (−0.8); tryptophan (−0.9); tyrosine (−1.3); proline (−1.6); histidine (−3.2); glutamate (−3.5); glutamine (−3.5); aspartate (−3.5); asparagine (−3.5); lysine (−3.9); and arginine (−4.5).

It is known in the art that certain amino acids may be substituted by other amino acids having a similar hydropathic index or score and still result in a protein with similar biological activity, *i.e.* still obtain a biological functionally equivalent protein. In making such changes, the substitution of amino acids whose hydropathic indices are within ± 2 is preferred, those which are within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred. It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity. U. S. Patent 4,554,101, incorporated herein by reference, states that the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with a biological property of the protein.

As detailed in U. S. Patent No. 4,554,101, the following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate ($+3.0 \pm 1$); glutamate ($+3.0 \pm 1$); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (−0.4); proline (-0.5 ± 1); alanine (−0.5); histidine (−0.5); cysteine (−1.0); methionine (−1.3); valine (−

1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); tryptophan (-3.4). It is understood that an amino acid can be substituted for another having a similar hydrophilicity value and still obtain a biologically equivalent, and in particular, an immunologically equivalent protein. In such changes, the substitution of amino acids whose hydrophilicity values are within ± 2 is preferred, those which are within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred.

As outlined above, amino acid substitutions are generally therefore based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. Exemplary substitutions that take various of the foregoing characteristics into consideration are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine and isoleucine.

Example 8 - Pesticidal Use of RNAi.

One method for disrupting transporter/channel protein function according to the subject invention is by the use double-stranded interfering RNA (RNAi), or RNA-mediated interference (RNAi). When RNAi corresponding to a sense and antisense sequence of a target mRNA is introduced into a cell, the targeted mRNA is degraded and protein translation of that message is stopped. Although not yet fully understood, the mechanism of this post-transcriptional gene silencing appears to be at least partially due to the generation of small RNA molecules, about 21 - 25 nucleotides in length, that correspond to (preferably at least 90% identity) the sense and antisense pieces of the RNAi introduced into the cell (Bass, B. L., 2000 "Double-stranded RNA as a template for gene silencing" *Cell* 101:235-238).

The specificity of this gene silencing mechanism is extremely high, blocking expression only of targeted genes, while leaving other genes unaffected (Chuang, C.-F. and E. M. Meyerowitz, 2000 "Specific and heritable genetic interference by double-stranded RNA in *Arabidopsis thaliana*" *Proc. Natl. Acad. Sci. USA* 97:4985-4990).

dsRNA (RNAi) typically comprises a polynucleotide sequence identical to a target gene (or fragment thereof) linked directly, or indirectly, to a polynucleotide sequence complementary to the sequence of the target gene (or fragment thereof). The dsRNA may comprise a

polynucleotide linker (stuffer) sequence of sufficient length to allow for the two polynucleotide sequences to fold over and hybridize to each other; however, a linker sequence is not necessary. The linker (stuffer) sequence is designed to separate the antisense and sense strands of RNAi significantly enough to limit the effects of steric hindrances and allow for the formation of dsRNA molecules.

RNA containing a nucleotide sequence identical to a fragment of the target gene is preferred for inhibition; however, RNA sequences with insertions, deletions, and point mutations relative to the target sequence can also be used for inhibition. Sequence identity may be optimized by sequence comparison and alignment algorithms known in the art (see Gribskov and Devereux, *Sequence Analysis Primer*, Stockton Press, 1991, and references cited therein) and calculating the percent difference between the nucleotide sequences by, for example, the Smith-Waterman algorithm as implemented in the BESTFIT software program using default parameters (e.g., University of Wisconsin Genetic Computing Group). Alternatively, the duplex region of the RNA may be defined functionally as a nucleotide sequence that is capable of hybridizing with a fragment of the target gene transcript.

RNA may be synthesized either *in vivo* or *in vitro*. Endogenous RNA polymerase of the cell may mediate transcription *in vivo*, or cloned RNA polymerase can be used for transcription *in vivo* or *in vitro*. Inhibition may be targeted by specific transcription in an organ, tissue, or cell type; stimulation of an environmental condition (e.g., infection, stress, temperature, chemical inducers); and/or engineering transcription at a developmental stage or age. The RNA strands may or may not be polyadenylated; the RNA strands may or may not be capable of being translated into a polypeptide by a cell's translational apparatus.

Preferably and most conveniently, RNAi can be targeted to an entire polynucleotide sequence of a gene set forth herein. Preferred RNAi molecules of the instant invention are highly homologous or identical to the polynucleotides shown in SEQ ID NO:1. The homology is preferably greater than 90% and is most preferably greater than 95%.

Fragments of genes can also be targeted. These fragments are typically in the approximate size range of about 20 nucleotides. Thus, targeted fragments are preferably at least about 15 nucleotides. In certain embodiments, the gene fragment targeted by the RNAi molecule is about 20-25 nucleotides in length. However, other size ranges can also be used. For example,

RNAi "fragments" of about 60 nucleotides with between 95 and 100% identity (to the target gene) can be used.

Genetic regulatory sequences, such as promoters, enhancers, and terminators, can be used in genetic constructs to practice the subject invention. Various constructs can be used to achieve
5 expression in specific plant tissues (by using root specific promoters, for example) and/or to target specific insect pest tissues (by using targeting elements or adjacent targeting sequences, for example).

In a specific embodiment of the subject invention, plant cells are genetically modified to produce at least one RNAi that is designed to be taken up by pests during feeding to block
10 expression (or the function of) of a target gene. As is known in the art, RNAi can target and reduce (and, in some cases, prevent) the translation of a specific gene product. RNAi can be used to reduce or prevent message translation in any tissue of the pest because of its ability to cross tissue and cellular boundaries. Thus, RNAi that is contacted with a pest by soaking, injection, or consumption of a food source will cross tissue and cellular boundaries. RNAi can
15 also be used as an epigenetic factor to prevent the proliferation of subsequent generations of pests.

Polynucleotide sequences disclosed herein can be used to identify conserved nucleotide motifs. Conserved nucleotide motifs strongly suggest that these sequences are functionally conserved. The use of these polynucleotides, and RNAi inhibitors thereof, is advantageous
20 because such RNAi can be designed to have broad RNAi specificity and are thus useful for controlling a large number of pests *in planta*.

Methods of the subject invention include the transformation of plant cells with genes or polynucleotides of the present invention, which can be used to produce RNAi in the plants. In one embodiment, the transformed plant or plant tissue can express RNAi molecules encoded by
25 the gene or polynucleotide sequence introduced into the plant. Other pesticidal constructs contemplated by the invention include antisense molecules specific to the polynucleotide sequences disclosed herein. The transformation of plants with genetic constructs disclosed herein can be accomplished using techniques well known to those skilled in the art and can involve modification of the gene(s) to optimize expression in the plant to be made resistant to

pests. Furthermore, it is known in the art that many tissues of the transgenic plants (such as the leaves, stems, fruit, or roots) can be targeted for transformation.

Example 9 - The efficacy of L-methionine as a *Manduca sexta* hornworm pesticide.

5 The effect of feeding L-methionine to *M. sexta* larvae as they progressed from the first through fifth instar stages was determined. Colonies of hornworms are commonly reared for research by well-accepted techniques that account for environmental conditions such as feed availability, daily photoperiod, temperature, *etc.* The caterpillars were obtained from Carolina Biological Supply Co. and were placed in a lighted 27°C incubator and fed hydrated
10 reconstituted Dry Tobacco Hornworm Medium Stock No. L-908D (Carolina Biological Supply Co., Burlington, NC) supplemented with or without (control) L-methionine supplementation (10% wt./wt. of dry components).

15 The effect of feeding L-Leucine or L-methionine to *M. sexta* larvae as they progressed from the first through fifth instar stages was studied. In one study, the caterpillars were placed in a 27°C incubator (photoperiod 16 hours light, 8 hours dark; relative humidity 60%) and fed a hydrated standard defined hornworm diet without (control) or with L-methionine supplementation (10% wt./wt. of dry components). Methionine supplementation prevented all the caterpillars from advancing in development, and killed 100% of the larvae within 3 days. None of the control larvae were killed -- indeed all the control larvae were healthy and proceeded
20 along their developmental course through at least the fifth instar stage (leading to the most crop-destructive stage). The results of this study are shown in Figure 1.

25 Table 3 reports the percentage treatment mortality of *Maduca* exposed to different concentrations of methionine over a 35 day period. Sample sizes of 20 neonate larvae per replicate (N=60 for each treatment; total N=480). Significance (*) was determined by one-way ANOVA using a Tukey Multiple Comparison at $P=0.05$, $df=16$, $q_{05,16,7} = 4.879$.

Table 3.		
Treatment	Treatment Mortality	Q
Control (0%) vs. 0.3% Methionine	0%	4.441
Control vs. 0.5%	63%	5.724*
Control vs. 1.0%	100%	6.662*
Control vs. 3.0%	100%	6.662*
Control vs. 5.0%	100%	6.662*
Control vs. 10.0%	100%	6.662*

Another study evaluated the *Manduca* larvae mortality on excised leaves of host plant (eggplant; *Solanum melongena*) that were sprayed with various concentrations of L-methionine. *Manduca Sexta* neonate larvae were initially reared on eggplants in Plexiglas cylinders. Larvae (total N=100) were exposed to eggplant leaves that were sprayed (then dried) with aqueous solutions of methionine. *Manduca* were 100% dead at all test doses, while the control (0% methionine) yielded zero percent mortality. Times to reach 100% mortality ranged from 2 days on leaves treated with 0.5% (w/w) methionine to 16 days on leaves treated with 0.06% (w/w) methionine. The results of this study are shown in Figure 2. In a further study, similar results were obtained with using intact plants growing in temperature controlled chambers. In the case of intact host plants, plant viability was not affected by L-methionine treatment, while the control plants were severely damaged by surviving larvae.

Example 10 - Efficacy of L-methionine as a *Manduca sexta* pesticide on host plant leaves using a Silwet-77 solution.

In one study, leaves of eggplant (*Solanum melongena*) were sprayed with a deionized water/Silwet-77 solution containing varying concentrations of methionine, then allowed to dry. The Silwet-77 was prepared according to the manufacture's instructions. *Manduca sexta* larvae (N=400; 80 per methionine dose) were then exposed to the eggplant leaves during 10 days, and surviving larvae were counted daily. The results of this study are shown in FIG 3.

Example 11 - Efficacy of L-methionine as a *Leptinotarsa decemlineata* pesticide.

Leaves of host eggplant were sprayed with deionized water containing varying concentrations of methionine, then dried. *Leptinotarsa decemlineata* larvae (N=20 per group) were exposed to the leaves, and surviving larvae were counted daily. The results of this experiment are shown in FIG 4. In this example, complete (100%) mortality was attained in 2 days with 0.47% methionine, compared to zero percent mortality with the control (0% methionine).

Example 12- Determination that L-methionine treatment for *L. decemlineata* or *M. sexta* does not adversely affect host plant fruit health, fruit yields or fruit weights.

Host eggplant (*Solanum melongena*) growing in fields were sprayed with deionized water containing varying concentrations of methionine, then dried. At all L-methionine concentrations, fruit yield mean numbers or mean weights were not significantly different from control (0% methionine) ($P \geq 0.05$; $N \geq 174$). The results of this experiment are shown in FIG. 5.

Example 13 - Assessment of efficacy of L-methionine as a mosquito pesticide.

Here, *Aedes aegypti* larvae entering the third or fourth instar stages (N=240; 40 per group) were placed in water containing various concentrations of L-methionine. In each solution, larvae fed *ad libitum* on commercial dried fish food applied to the water. The results of this study are shown in FIG 6.

Example 14- Further Assessment of Efficacy of L-methionine as a mosquito pesticide.

Another experiment assessed the specificity of L-methionine compared to L-proline on *Aedes aegypti* survival. Mosquito larvae (third or fourth instar; N=240) were placed in water containing various concentrations of L-proline or L-methionine, and fed *ad libitum* with commercial dried fish food applied to the water. There was no effect of L-proline compared to 0% methionine control ($P \geq 0.05$) at all test days. However, 0.7% methionine killed 100% larvae with LD50 ~ 0.3% methionine on day 3. Similar data were obtained for day 2, which yielded LD50 ~ 0.5% methionine. See Figure 7.

Example 15 - Pests with Alkaline Gut Compartments.

Table 4 provides an illustrative, non-exhaustive, list of pests having alkaline gut compartments.

Table 4 provides an illustrative, non-exhaustive, list of pests having alkaline gut compartments.

Table 4 - The pH in various parts of the gut of selected insects

Insect	Foregut	Midgut	Hindgut	Reference
<u>Orthoptera</u>				
<i>Photalliotus nebrascensis</i> grasshopper (Acrididae)	6.03	7.12	6.11	(1)
<i>S. gregaria</i> desert locust		5.3		(5)
<i>Gryllus rubens</i> field cricket (Gryllidae)	5.8-6.0	7.4-7.6	7.6-7.8 (ant hindgut)	(7)
<i>Gryllus bimaculatus</i> field cricket (Gryllidae)	5.84 (crop)	8.07 (ventriculus)	8.50 7.59 (ileum)(rectum)	(20)
<i>Leucophaea madeirae</i> cockroach (Blattidae)	9.5 (posterior midgut)			(9)
<u>Coleoptera</u>				
<i>Popillia japonica</i> larvae Japanese beetle (Scarabaeidae)	8.5			(2)
<i>Exomala orientalis</i> larvae Oriental beetle (Scarabaeidae)	8.5-9.0			(2)
<i>Rhizotrogus majalis</i> larvae European chafer (Scarabaeidae)	9.0-9.5			(2)
<i>Maladera castanea</i> larvae Asiatic garden beetle (Scarabaeidae)	8.5			(2)

Table 4 (continued)
Insect

	Foregut	Midgut	Hindgut	Reference
<u>Coleoptera</u>				
<i>Lichnanthe vulpina</i> larvae Cranberry root grub (Scarabaeidae)	8.5			(2)
<i>Phyllophaga anxia</i> larvae Phyllophaga white grub (Scarabaeidae)	8.5-9.0			(2)
<i>Oryctes nasicornis</i> (Scarabaeidae)	12.2			(15)
<u>Lepidoptera</u>				
<i>Agrotis ipsilon</i> black cutworm (Noctuidae)	8.5-9.0			(2)
<i>Manduca sexta</i> tobacco hornworm (Sphingidae)	9.5-9.7			(5)
<i>Manduca sexta</i>	6.4 8.2 7.2 7.1	apical folds of ant. midgut basal folds of ant. midgut Apical folds of post midgut Basal folds of post midgut		(6)
<u>Diptera</u>				
mosquito larva	~10			(14)
<i>Simulium vitatum</i> Blackfly (Simuliidae)	11.4			(16)

Table 4 (continued)
Insect

	Foregut	Midgut	Hindgut	Reference
	<u>Diptera</u>			
<i>Tipula abdominalis</i> crane fly (Tipulidae)	11.6			(17)
<i>Lucilia cuprina</i> larva blowfly (Calliphoridae)	7.4-8 anterior midgut	3.3 middle midgut	7.4-8 posterior midgut	(18)
	<u>Isoptera</u>			
termites	>	10 (ant. midgut)		(12)

References; Many additional references can be found in Berenbaum (1980).

- (1) Barbehenn, R.V., M.M. Martin, and A.E. Hagerman. 1996. Reassessment of the roles of the peritrophic envelope and hydrolysis in protecting polyphagous grasshoppers from ingested hydrolyzable tannins. J. Chem. Ecol. 22: 1911-1929.
- (2) Broadway, R.M. and M.G. Villani. 1995. Does host range influence susceptibility of herbivorous insects to non-host plant proteinase inhibitors? Entomol. Exp. et Appl. 76: 303-312.
- (3) Murdock, L.L., G. Brookhart, P.E. Dunn, D. E. Foard, S. Kelley, L. Kitch, R.E. Shade, R.H. Shukle, and J.L. Wolfson. 1987. Cysteine digestive proteinases in Coleoptera. Comp. Biochem. Physiol. 87B: 783-787.
- (4) Evans, W.A.L., and D.W. Payne. 1964. Carbohydases of the alimentary tract of the desert locust, *Schistocerca gregaria* Forsk. J. Insect Physiol. 10: 657-674.
- (5) Martin, J.S., M.M. Martin, and E.A. Bernays. 1987. Failure of tannic acid to inhibit digestion or reduce digestibility of plant protein in gut fluids of insect herbivores: Implications for theories of plant defense. J. Chem. Ecol. 13: 605-621.
- (6) Dow, J.A.T., and M.J. O'Donnell. 1990. Reversible alkalization by *Manduca sexta* midgut. J. Exp. Biol. 150: 247-256.
- (7) Thomas, K.K. and J. L. Nation. 1984. Protease, amylase and lipase activities in the midgut and hindgut of the cricket, *Gryllus rubens* and mole cricket, *Scapteriscus acletus*. Comp.
- (8) O'Riordan, A.M. 1969. Electrolyte movement in the isolated midgut of the cockroach (*Periplaneta americana*). J. Exp. Biol. 51: 699-714.

- (9) Engelmann, F., and P.M. Geraerts. 1980. The proteases and the protease inhibitor in the midgut of *Leucophaea maderae*. J. Insect Physiol. 26: 703-710.
- (10) Martin, M.M., J.J. Kukor, J.S. Martin, D.L. Lawson, and R.W. Merritt. 1981a. Digestive enzymes of larvae of three species of caddisflies (Trichoptera). Insect Biochem. 11: 501-505.
- (11) Martin, M.M., J.S. Martin, J.J. Kukor, and R.W. Merritt. 1981b. The digestive enzymes of detritus-feeding stonefly nymphs (Plecoptera; Pteronarcyidae). Can. J. Zool. 59: 1947-1951.
- (12) Bignell, D.E., and J.M. Anderson. 1980. Determination of pH and oxygen status in the guts of lower and higher termites. J. Insect Physiol. 26: 183-188.
- (13) Mishra, S.C., and P.K. Sen-Sarma. 1981. Hydrogen ion concentration in the digestive tract of three species of Indian termites. Entomon. 6: 131-134.
- (14) Dadd, R.H. 1975. Alkalinity within the midgut of mosquito larvae with alkaline-active digestive enzymes. J. Insect Physiol. 21: 1847-1853.
- (15) Bayon, C. 1980. Volatile fatty acids and methane production in relation to anaerobic carbohydrate fermentation in *Oryctes nasicornis* larvae (Coleoptera: Scarabaeidae). J. Insect Physiol. 26: 819-828.
- (16) Undeen (1979)
- (17) Martin et al. (1980)
- (18) Waterhouse, D.F., and B. Stay. 1955. Functional differentiation in the midgut epithelium of blowfly larvae as revealed by histochemical tests. Aust. J. Biol. Sci. 8: 253-277.
- (19) Krishna, S.S., and K.N. Saxena. 1962. Digestion and absorption of food in *Tribolium castaneum* (Herbst.). Physiol. Zool. 35:66-78.
- (20) Teo, L.H. 1997. Tryptic and chymotryptic activities in different parts of the gut of the field cricket *Gryllus bimaculatus*

Example 16 - Insertion of Toxin Genes Into Plants.

One aspect of the subject invention is the transformation of plants with genes encoding the pesticidal compounds of the present invention. The transformed plants are resistant to attack by the target insect pest.

5 Genes encoding pesticidal compounds, as disclosed herein, can be inserted into plant cells using a variety of techniques that are well known in the art. For example, a large number of cloning vectors comprising a replication system in *E. coli* and a marker that permits selection of the transformed cells are available for transforming higher plants, e.g., pBR322, pUC series, M13mp series, pACYC184, etc. Accordingly, the sequence encoding the pesticidal peptide can
10 be inserted into the vector at a suitable restriction site. The resulting plasmid can be used for transformation into *E. coli*. The *E. coli* cells can be cultivated in a suitable nutrient medium, then harvested and lysed. The plasmid can be recovered. Sequence analysis, restriction analysis, electrophoresis, and other biochemical and/or molecular biological methods can be generally carried out as methods of analysis. After each manipulation, the DNA sequence used
15 can be cleaved and joined to the next DNA sequence. Each plasmid sequence can be cloned in the same or other plasmids. Depending on the method of inserting desired genes into the plant, other DNA sequences may be necessary. If, for example, the Ti plasmid (the tumor-inducing plasmid of the plant-pathogenic bacterium *Agrobacterium tumefaciens*) or Ri plasmid (the root-inducing plasmid of *Agrobacterium rhizogenes*) can be used for the transformation of the plant
20 cell, then at least the right border, but often the right and the left border of the Ti or Ri plasmid T-DNA ("Transferred DNA"), must be joined as the flanking region of the genes to be inserted.

A large number of techniques are available for inserting DNA into a plant host cell. These techniques include transformation with T-DNA using *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes* as transformation agent, fusion, injection, biolistics (microparticle
25 bombardment), or electroporation as well as other possible methods.

One of the most widely used approaches for the introduction of DNA into plant cells exploits the natural DNA-transferring properties of *Agrobacterium tumefaciens* and *Agrobacterium rhizogenes*, the two species which cause crown gall and hairy root. Their ability to cause disease depends on the presence of large plasmids, in excess of 100 kb, which
30 are referred to as the Ti and Ri plasmids, respectively.

A region referred to as the T-DNA ("Transferred DNA") is transferred from an infecting *Agrobacterium* cell into the nucleus of the plant cell, where it is integrated into the plant genome. The use of T-DNA for the transformation of plant cells has been intensively researched and sufficiently described in Eur. Pat. Appl. No. EP 120 516; Hockema (1985); An
5 *et al.*, 1985, Herrera-Estrella *et al.*, (1983), Bevan *et al.*, (1983), and Klee *et al.*, (1985). Transfer of the T-DNA depends on a set of genes called *vir* if they are on the Ti plasmid, or *chv* if they are on the chromosome. These genes are induced in response to various compounds in exudates from wounded plants. The T-DNA itself is flanked by repeated sequences of around 25 base pairs, called border repeats (or left and right borders). The T-DNA contains a group of
10 genes referred to as the *onc* genes, which are responsible for the oncogenicity of the T-DNA.

The use of *Agrobacterium* in the genetic manipulation of plants involves the insertion of foreign DNA into the T-DNA of a bacterial cell and subsequent transfer of the DNA by the transformed bacterium into the plant. As long as the necessary proteins are provided by the bacterium, any sequences flanked by the T-DNA border repeats can be transferred into the
15 recipient plant cell genome. The Ti plasmids are too large to manipulate directly, but this problem can be circumvented by using cointegrative and binary systems.

The two main components of a cointegrative system are a Ti plasmid that has typically been modified by the replacement of material between the border repeats (including the *onc* sequences) by pBR322; and an intermediate vector, which is a modified pBR322 containing an
20 extra marker, such as kanamycin resistance. The gene to be introduced into the target plant is first cloned into the intermediate vector, and this construct is then introduced into *Agrobacterium* containing the Ti vector. The pBR322-based plasmid cannot replicate efficiently inside *Agrobacterium*, so selection for kanamycin resistance identifies those *Agrobacterium* cells where the pBR322-based intermediate plasmid has been integrated by
25 homologous recombination into the Ti plasmid. Because the recombination is homologous, it will take place across the pBR322 sequences and therefore result in integration between the border repeats.

The need for cointegration of the plasmids can be circumvented by use of a binary vector, such as pBin19, a small plasmid containing a pair of left and right borders. The *lacZ*
30 region, located within the borders, facilitates insertion and detection of DNA. A neomycin

phosphotransferase gene, typically modified for expression in plants by addition of nopaline synthase expression sequences, is also present within the borders. Outside the left and right borders, there is typically a kanamycin resistance gene that will function in prokaryotes and a broad host-range origin derived from the plasmid pRK252. The proteins that catalyze transfer of the T-DNA into the host plant do not have to be *cis*-encoded (*i.e.*, do not have to be encoded by the same molecule). Therefore, if the binary vector is introduced into *Agrobacterium* that already contains a resident Ti plasmid, the resident plasmid can provide all the functions needed to transfer into a plant nucleus the DNA between the borders of the binary vector. Other, more sophisticated binary vectors, are also known in the art, for example pROK1. These vectors typically have plant promoters incorporated to drive expression. Others have *cos* sites to allow packaging into lambda phage heads.

When the correct sequences have been incorporated into a vector (whether binary or cointegrative), the vector must then be transferred to an *Agrobacterium* strain carrying an appropriate Ti plasmid. This is usually accomplished either by electroporation with naked DNA or by a triparental mating involving the *Agrobacterium* strain, an *E. coli* strain containing the vector to be transferred, and an *E. coli* strain with a plasmid capable of mobilizing the binary or intermediate vector into *Agrobacterium*.

Once the binary vector of the cointegrative vector has been introduced into a suitable *Agrobacterium* strain (and cointegration has occurred), the next stage is to permit the *Agrobacterium* to infect plant cells. Various methods exist, including inoculation of intact plants with *Agrobacterium* cultures by injection, but the most widely used is to incubate discs cut from leaves of the target plant with an *Agrobacterium* culture. The bacterium will attack cells around the edge of the wounded leaf disc and transfer its T-DNA back into them. The leaf discs are then transferred to a suitable medium to select for transformation. The neomycin phosphotransferase gene is widely used, conferring resistance to aminoglycoside antibiotics, such as neomycin, kanamycin, and G518. On a suitable selective medium, shoots form around the edges of the treated leaf discs. The shoots can then be regenerated into intact plants. See Howe, *Gene Cloning and Manipulation* (1995).

The transformed cells are regenerated into morphologically normal plants in the usual manner. If a transformation event involves a germ line cell, then the inserted DNA and

corresponding phenotypic trait(s) will be transmitted to progeny plants. Such plants can be grown in the normal manner and crossed with plants that have the same transformed hereditary factors or other hereditary factors. The resulting hybrid individuals have the corresponding phenotypic properties.

5

It should be understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200